

HELPING PEOPLE HELP THEMSELVES

Evidence about supporting self management

October 2010

Identify Innovate Demonstrate Encourage

KEY MESSAGES

An increasing number of people are living with long term health conditions, which they manage most of the time by themselves. Helping people to better care for themselves can improve their physical and mental wellbeing and change how they use services. There is good evidence that supporting self management works.

What is self management?

As the population ages and our lifestyles and habits change, more and more people are living with long term conditions that cannot currently be cured. In the UK more than 17 million adults have a long term condition such as diabetes, asthma, heart failure, arthritis dementia or depression.

Health and social services support people with long term conditions by providing equipment, specialist staff and medicines to control symptoms. In fact, it is estimated that about two thirds of all healthcare resources are spent supporting people with long term conditions.

Even so, 80% to 90% of all care for people with long term conditions is undertaken by patients themselves and their families. This 'self management' or 'self care' includes eating well, exercising, taking medicines, keeping in good mental health, watching for changes, coping if symptoms worsen and knowing when to seek professional help.

Supporting self management is about educating people about their condition and care and motivating them to care for themselves better. Planners and practitioners increasingly recognise that supporting people to self manage their condition can improve symptoms and clinical outcomes and may change how people use health services.

Does self management work?

This rapid review compiles research evidence about supporting self management. Based on the results of almost 600 studies published in the UK and internationally, there is evidence that **supporting self management can improve people's quality of life, clinical outcomes and health service use.**

Some studies and reviews suggest that the evidence for supporting self management is only moderate but this is because a wide range of things are described as 'self management support' and some may be more effective than others. Past reviews have combined initiatives that focus solely on information provision with interventions that more actively target behaviour change and self efficacy. However these varying interventions may have different outcomes so combining them could dilute the findings.

Although some studies suggest supporting self management reduces the use of health services, this is not yet generalisable. Focusing on reducing healthcare use and costs may also be too simplistic. It is more likely that **patterns of service use change** rather than reduce overall. For example, people may engage more frequently with a practice nurse, telephone health coach or peers, but less with hospital services. The aim is not to reduce contact overall, but rather to support a different pattern of contact which may lead to fewer crises and inpatient admissions.

What type of support works?

There are a wide range of initiatives to support self management including information leaflets, online peer support, one to one counselling, group education sessions, telephone coaching, using technology to monitor symptoms and psychological behaviour change interventions.

Initiatives can be categorised along a continuum, with passive information provision about people's condition and 'technical skills' at one end of the scale and initiatives that more actively seek to support behaviour change and increase self efficacy at the other end of the continuum.

All of the different types of support are important components of the jigsaw needed to encourage self management, but **information provision alone is unlikely to be sufficient to motivate behaviour change** and improve clinical outcomes. More active goal setting and behavioural change interventions are also necessary.

Different clinical conditions may require varying approaches to support self management. For instance, because of the nature of conditions such as diabetes, there is a role for structured patient education focused on technical or clinical information about diet, exercise and medication. For conditions such as depression on the other hand, there may be fewer 'technical' changes that people can make so the focus is instead on cognitive and behavioural interventions. Such interventions may be equally valuable for people with diabetes and other conditions where the focus to date has been mainly on providing 'technical' information, but this remains an emerging field of knowledge.

What else do we need to know?

There is evidence that several general principles are important when supporting self management, including **promoting** healthy lifestyles and **educating** people about their conditions and how to self manage; **involving** people in decision making; **motivating** people to self manage using targeted approaches and structured support; helping people **monitor** their symptoms and know when to take appropriate action; developing care plans as a **partnership** between service users and professionals; setting **goals** and following up on the extent to which these are achieved over time; helping people manage the social, emotional and physical **impacts** of their conditions; proactive **follow up** and providing opportunities to share with and learn from other **service users**.

However, knowledge about the **best strategies** to support behaviour change and put these principles into practice is still being developed. Research suggests that in order to change behaviour, people need to want to change and 'stage of change' models have been found useful in developing successful programmes.

Another core component is **supporting clinicians** to work with people to improve their motivation to change. It is important to understand the skills that clinicians need to help people make changes. There may also be a need to fundamentally change the way patients and practitioners see their roles, but to date evidence about this is limited.

To conclude, research suggests that supporting self management works, and can have real impacts on how people think, feel and act. The challenge is to explore the best ways to support self management and to help service users, clinicians and managers make this a reality.

CONTENTS

What is self management?	4	
Does self management work?	5	
Impact on self efficacy	5	
Impact on self care behaviour	6	
Impact on quality of life	7	
Impact on clinical outcomes	8	
Impact on health service use	10	
What type of support works?	14	
Providing information	15	
Providing support	15	
Behaviour change approaches	16	
What else do we need to know?	17	
Changing behaviour	17	
Supporting clinicians	17	
Summary	18	
Self management works	18	
Proactive strategies work best	18	
Evidence is developing	19	
Appendix 1	Review methods	20
Appendix 2	Research details	21
References		29

WHAT IS SELF MANAGEMENT?

Health services in Britain are facing an enormous challenge. The population is growing in size and age and people are more likely to suffer from long term illnesses that require ongoing care.¹ There is a need to change the way systems work and this includes helping people to help themselves.

Long term conditions are leading causes of death and disability worldwide.² In Britain, the Department of Health estimates that 17.5 million adults may be living with one or more long term conditions.³ 60% of adults report having at least one long term condition⁴ and this figure continues to grow due to an aging population and escalating risk factors such as obesity.

Medical advances mean that people with illnesses such as heart disease and stroke now receive treatments that enable them to live longer and to enjoy a higher quality of life than would previously have been possible, but this also brings challenges. Health and social care services provide help when symptoms worsen, but most of the time people and their families are responsible for making decisions and influencing their wellbeing. In the UK about 80% of GP consultations, 60% of days spent in hospital and two thirds of emergency admissions are related to long term conditions,⁵ but more than 80% of the care for people with long term illnesses is undertaken by the patient themselves or their carers.⁶ This is known as self management.

The Department of Health defines self management as

“the actions individuals and carers take for themselves, their children, their families and others to stay fit and maintain good physical and mental health; meet social and psychological needs; prevent illness or accidents; care for minor ailments and long term conditions; and maintain health and wellbeing after an acute illness or discharge from hospital.”⁷

Supporting self management involves providing information and encouragement to help people maintain greater control by understanding their condition and being able to monitor and take appropriate action. Interventions to support self management can be used at different points of the health continuum, from those who do not have a long term condition through to those who are living with severe and multiple long term conditions.

Health and social care services can support people to self manage their conditions by encouraging engagement in decision making, educating people about their condition and care, motivating people to adopt healthy behaviours and helping them know when and how to seek help from professionals.⁸

Over the past decade there has been an increasing focus in the UK and around the world on supporting self management. Many strategies have been tested, with varying success, including providing accessible information, communication skills training for service users and professionals, self management skills training facilitated by lay people, nurse led telephone support, self monitoring of clinical symptoms, and even text messages, computer forums and video games. This rapid review compiles evidence about the effects of supporting self management on people’s quality of life, clinical outcomes and health service use.

The review methods are summarised in Appendix 1. In brief, two reviewers searched more than 10 bibliographic databases for research evidence published up until September 2010. More than 100,000 reports were scanned and the findings from over 550 high quality studies are included in the review.

The review does not aim to be exhaustive but instead provides a **rapid and easy to use** compilation of up to date evidence. First, we explore whether supporting self management improves outcomes, then we examine the most effective types of support and areas in need of further development.

DOES SELF MANAGEMENT WORK?

Evidence suggests that supporting self management works. **Supporting people to look after themselves can improve their motivation, the extent to which they eat well and exercise, their symptoms and clinical outcomes and can even change how they use health services.**

Interventions to encourage and support self management vary considerably in their aims, approach, content, delivery, duration and target group.⁹ Therefore it would be misleading to refer to ‘self management initiatives’ as an integrated whole. This section describes evidence about the impact of supporting self management on people’s attitudes, behaviours and outcomes, but it is important to emphasise from the outset that these results vary according to the type of support provided. The next section explores differences between various approaches.

Impact on self efficacy

How people think and feel about their condition can have a big impact on their health behaviours and outcomes.^{10,11,12,13,14}

“Self efficacy refers to an individual’s belief in their capacity to successfully learn and perform a specific behaviour. A strong sense of self efficacy leads to a feeling of control, and willingness to take on and persist with new and difficult tasks. When applied to health, this theory suggests that patients are empowered and motivated to manage their health problems when they feel confident in their ability to achieve this goal.”¹⁵

There is evidence that improved self efficacy is correlated with improved health behaviours and clinical outcomes so it is valid to examine the impact of self management support on self efficacy as a proxy for other outcomes.^{16,17,18,19,20,21,22,23,24,25,26,27,28}

A literature review found that involving people in ongoing health decision making and self management can increase patient and family responsibility for the delivery of care, and help people adapt care regimens to their own lifestyles.²⁹

However, the best strategies to encourage self efficacy remain uncertain. A trial in the US examined the impact of empowerment and motivation on self management behaviours. Those who felt more empowered and active were most likely to self manage effectively. However specific self management education programmes did not seem to make people more activated. The authors concluded that more research is needed to examine the best ways to support self management in order to increase empowerment and motivation.³⁰ Others suggest that it is important to understand why people wish to receive self management support in order to target assistance to their needs.³¹

A number of self management interventions focus on confidence building and providing service users with the knowledge and skills to set personal goals and develop effective strategies for achieving them.^{32,33,34,35,36,37,38,39,40,41}

For instance, UK studies suggest that people with arthritis taking part in self management programmes feel more confident in their ability to manage and control their symptoms, feel less anxious about their disease, and may visit the doctor less frequently.^{42,43,44,45} Similar studies are available for people with heart disease, lung disease, diabetes, asthma, stroke and many other conditions.^{46,47,48,49,50,51}

Some suggest that supporting self management can help people move through **stages of change** (transtheoretical model), becoming more motivated to alter their behaviours and sustain this long term.^{52,53,54,55,56} Others have found that the effectiveness of self management support varies depending on people’s stage of change.^{57,58,59,60,61,62,63,64,65,66,67,68}

Box 1: examples of targeting self efficacy

A small randomised trial in Taiwan investigated the effectiveness of an empowerment programme in 50 people with end stage renal disease. The programme included identifying problem areas for self management, exploring emotions associated with these problems, developing a set of goals and strategies to overcome problems and achieve goals, creating and implementing behavioural change plans and stress management. There were improvements in empowerment, self care, self efficacy and depression.⁶⁹

Similarly, a trial in China examined layperson led self management education for 954 people with hypertension, heart disease, chronic lung disease, arthritis, stroke, or diabetes.⁷⁰ There were improvements in self care behaviour, self efficacy, and health status and reduced hospitalisations six months after the course. Integrating the delivery model into community organisations and working with service users as lay educators were key success factors.

Impact on self care behaviour

There is evidence that self management support can improve people's knowledge about their condition and care, how they feel about their condition, their ability to cope day to day and self management behaviours.^{71,72,73,74,75,76,77,78,79,80,81}

For instance, a meta analysis of 82 studies found that self management education improved knowledge, self care behaviour, and metabolic control in adults with diabetes.⁸²

Another review of randomised trials found that self management education increases participants' self efficacy, knowledge, symptom management, use of self management behaviours and aspects of health status. The effects for children, young adults, and carers remains uncertain.⁸³

Box 2: examples of impact on behaviour

Most of the available research focuses on the impact of group education sessions. For example, a five year randomised trial with more than 1000 people in the UK found that self management courses were associated with improved healthy behaviours, coping, communication with physicians and self reported health status and fewer days in hospital.⁸⁴

A comparison of routinely collected data for 15,190 older people with diabetes found that those who had attended self management education were more likely to self monitor blood glucose levels, take appropriate medications and go for regular eye examinations.⁸⁵

Benefits from other types of support have also been documented. For instance, researchers in the US found that sending weekly educational newsletters improved self management behaviours among African Americans with diabetes.⁸⁶

Research has explored the perceived barriers and facilitators for lifestyle modification. For example a qualitative study with people with COPD found that the most successful self management support requires a multifaceted approach incorporating strategies to show how to change behaviour. This and other studies suggest that self management support that includes behaviour change strategies and problem solving skills is more likely to lead to improved self efficacy and lifestyle modification.^{87,88,89,90}

Programmes which combine multiple support strategies often work best. A randomised trial in New Zealand included a diary for recording daily weights and goals, attendance at a specialist clinic, and three self management education sessions for people with heart failure. After one year, those taking part in the intervention were more likely to have better self management knowledge and behaviours compared to the usual care group. Patients who did not adopt self management strategies had a greater chance of death or readmission to hospital.⁹¹

However, other studies have found no difference between self management support and usual care regarding self efficacy, behaviours and awareness.⁹²

Most studies that find no benefits are small, perhaps non randomised and often focus on less structured forms of self management support. This re-emphasises that **not all forms of self management support are homogenous**, and nor are their outcomes.

Impact on quality of life

Research suggests that supporting self management can improve quality of life.^{93,94,95,96,97,98,99} A number of studies have found a link between self management education, self care behaviours and psychological outcomes, stress, coping or quality of life.^{100,101,102,103,104,105,106,107,108}

There is evidence that supporting self management results in both improved confidence to self manage and improved quality of life^{109,110,111,112} and that self efficacy or perceived control is correlated with improved quality of life.^{113,114,115}

Box 3: examples of interlinkages

There is a close relationship between self management attitudes, behaviours and quality of life. For instance, trials in China found that self management education was associated with improved self care behaviours which in turn were associated with improvements in quality of life, clinical outcomes and healthcare resource use.^{116,117}

Another randomised trial in the US assessed a self management programme for older people with deteriorating eyesight. The programme focused on enhancing problem solving skills and resulted in improved mood and reduced emotional distress, which in turn was associated with increased self efficacy. Functional status and quality of life also improved, especially for those who were initially depressed. This suggests a link between active self management education, self management behaviours and quality of life.¹¹⁸

Similarly, a randomised trial in the UK found that a self management planning intervention was associated with improved self care for people with angina, including better diet and more exercise. The programme was also associated with improved anxiety and quality of life.¹¹⁹

There are some conflicting findings. A Cochrane review found that self management education by lay educators may lead to **short term improvements** in self efficacy, self rated health status and self care behaviours but no improvements in quality of life.¹²⁰

Likewise, a randomised trial with 131 people with asthma in Switzerland found that self management education helped to improve self management skills, self efficacy and people's knowledge of their condition and care. However there was no improvement in quality of life or health outcomes.¹²¹

Another small trial in the UK found that supporting self management improved self efficacy and self care in people with stroke but did not improve quality of life.¹²²

Studies that have not found any effect tend to be small or to use less proactive strategies to support self management, such as written information or short education sessions focusing on providing technical information.

Impact on clinical outcomes

Overall, the evidence suggests that there is likely to be a relationship between supporting self management and clinical outcomes.^{123,124,125,126,127,128,129,130,131,132}

Some studies suggest direct relationships between supporting self management and improved clinical outcomes.^{133,134,135,136,137,138,139,140} Other studies show a correlation, but not necessarily a causal relationship.^{141,142,143,144,145,146}

Various types of self management support have been found to improve physical symptoms or clinical outcomes in people with arthritis, asthma, diabetes, hypertension, heart disease, heart failure, stroke, cancer and other conditions, at least in the short-term.^{147,148,149,150,151,152,153}

For instance, a systematic review collated 71 trials of self management education for people with long term conditions. The reviewers found that people with diabetes participating in self management education had improved glycaemic control and blood pressure. People with asthma experienced fewer attacks after self management education. Arthritis self management education programmes had no significant effects. The authors concluded that self management education programmes may have small to moderate effects for people with selected long term conditions. They noted, however, that study methods varied widely and were not optimal.¹⁵⁴

Box 4: examples of clinical improvements

A meta analysis assessed the effectiveness of self management education in children and adolescents with asthma. The reviewers included 32 randomised trials with 3706 children aged between 2 and 18 years. Self management education was associated with improved lung function and self-efficacy and reduced absenteeism from school, days of restricted activity, and visits to A&E. Programmes based on peak flow monitoring, targeted interventions, and interventions with children with severe asthma had the most effect on clinical outcomes.¹⁵⁵

Similarly, a cohort study in the US examined the relationship between self management education, self management behaviours and clinical outcomes for 428 people with diabetes. Education was associated with improved self care and more controlled blood glucose.¹⁵⁶ People in the early stages of illness or with less severe illness were more likely to benefit, and this finding has been replicated in studies of people with other conditions.^{157,158}

Another study in the US found that active self management education was associated with improved **empowerment** and engagement between people with diabetes and clinicians. This in turn resulted in improved glycaemic control.¹⁵⁹

A review of self management education in arthritis found reductions in pain and fatigue, improved activity levels, aerobic capacity and exercise endurance, reduced levels of disability and functional limitations, and improved self-reported health status.¹⁶⁰ Numerous similar studies are available about measures of pain and quality of life.^{161,162}

Studies have also assessed interventions involving the **family members** of people with long term conditions.¹⁶³ For instance a randomised trial with children with diabetes found that a family teamwork intervention prevented deterioration in glycaemic control.¹⁶⁴

There are exceptions to these positive trends.¹⁶⁵ Some studies suggest no improvements in clinical outcomes. For instance, a randomised trial in London evaluated a card and letter encouraging people to prepare questions to ask at a hospital consultation. Half of those sent a card said they got more out of their consultation as a result, but there were no significant differences in outcomes.¹⁶⁶ Another randomised trial with 131 people with asthma found that improved self management skills and self efficacy did not result in health improvements.¹⁶⁷

Others suggest that any improvements are **short lived**. A systematic review of 31 studies found that self management education for people with diabetes improves clinical outcomes in the short term. The longer the self management intervention, the longer the effect tends to last. Any benefits tend to decrease 1-3 months after the intervention ceases, suggesting that learned behaviours change over time.¹⁶⁸

As with other outcomes, the **level of engagement** and activity may be an important success factor for improving clinical outcomes. A randomised trial in Argentina compared self management education aimed at improving communication skills and empowerment versus a compliance based relationship between patients and professionals. The study found that the self management approach was associated with improved clinical outcomes; in this case increased blood pressure control.¹⁶⁹

However, some suggest research in this field is methodologically flawed. A systematic review of self management education for people with heart failure found that while there were small improvements overall in mortality and healthcare resource use, most studies were small and suffered from biases or methodological design flaws.¹⁷⁰

Impact on service use

There is evidence from the UK and elsewhere that self management support can alter the **pattern of healthcare service use** and subsequent healthcare costs, though the evidence is more varied than for clinical outcomes.^{171,172,173,174,175,176,177,178,179,180}

It has been suggested that self management support programmes may reduce visits to health services by up to 80%.^{181,182,183,184,185,186} Benefits may remain over the long term^{187,188} and have been found throughout the world.¹⁸⁹

Box 5: examples of impact on resources

Evidence about health service use tends to focus on complex mixed method approaches to support self management or group education courses.

As an example of a mixed method approach, a randomised trial in the UK assigned 203 people with ulcerative colitis undergoing hospital follow up to patient centred self management training and follow up on request, or usual care and routine follow up. Self management training was associated with faster access to treatment when needed, reduced hospital visits (0.9 versus 2.9 per patient per year), and fewer GP visits (0.3 versus 0.9 per patient per year).^{190,191} A success factor was giving service users control over access to care when they felt they needed it.

Another randomised trial in 19 hospitals in northwest England examined whether a whole systems approach to self management was cost effective among 700 people. Consultants were trained to provide a patient centred approach to care and patient guidebooks were developed in partnership with service users. All patients prepared written self management plans. After one year, the self management group had fewer hospital visits and patients felt more able to cope with their condition. Cost effectiveness analyses favoured self management over usual care.¹⁹²

Trials and descriptive studies in the UK suggest that group education sessions such as the Expert Patient Programme have the potential to reduce hospital admissions and days spent in hospital.^{193,194} However evaluations to date have been equivocal about the impact on healthcare resource use.¹⁹⁵

A survey with 1000 people who had attended an Expert Patient Programme in England found that people reported improved self management, reduced medication use and fewer unscheduled visits to the GP and A&E,¹⁹⁶ but administrative data were not available to confirm this.

Evidence is available about similar types of self management courses from around the world. A literature review found that structured self management education programmes improve self care behaviours and can reduce healthcare resource use and expenditure. Potential benefits include fewer A&E visits, fewer hospitalisations, and reduced days spent in hospital.¹⁹⁷ Results hold in resource poor settings too.¹⁹⁸

A meta analysis found that self management education programmes improved knowledge of diabetes and clinical outcomes and reduced medication use. The authors concluded that for every five people attending a group based education programme, one person would be expected to reduce diabetes medication and this would impact on overall costs.¹⁹⁹

Another example is a randomised trial in seven hospitals in Canada, which evaluated self management education among people with moderate to severe chronic obstructive pulmonary disease hospitalised within the past year. The programme involved weekly visits by health professionals over two months, with monthly telephone follow up. Self management education was associated with 40% fewer hospital visits for chronic obstructive pulmonary disease and 57% fewer hospital admissions for other problems. Emergency department visits reduced by 41% and unscheduled physician visits by 60%.²⁰⁰

Similarly, a trial in the US assessed seven weekly sessions of peer facilitated self management education, each of 2.5 hours duration, provided in community settings with groups of 15 to 20 participants. The programme reduced emergency department and outpatient visits, improved health behaviours, reduced symptoms, and improved health status. Reductions in service use and emotional distress were evident two years after the programme.²⁰¹

Another randomised controlled trial in Norway assessed self management education for people with asthma. The intervention included two education sessions plus two follow ups by nurses or physiotherapists. At one year follow up those taking part in self management education had reduced visits to GPs, reduced overall costs and improved quality of life.²⁰²

A randomised trial with children with asthma examined self management education using interactive multimedia tools. The initiative was associated with increased asthma knowledge among children and caregivers, decreased asthma symptom days, fewer emergency department visits, and lower average daily doses of inhaled corticosteroids. Increased asthma self management knowledge and behaviours were associated with fewer urgent physician visits and less frequent use of quick relief medicines.²⁰³

Numerous trials are available suggesting a link between self management education and reduced or altered patterns of health service use.^{204,205,206,207,208, 209} But we need to be cautious when interpreting the findings because some research suggests correlations rather than direct links. Many studies have found that supporting self management improves self care knowledge or behaviours and reduces hospitalisations but they have not proven that self care causes reduced resource use.^{210,211,212,213,214}

The mechanism by which outcomes change remains uncertain. A randomised trial in the US for people with chronic depression and post traumatic stress disorder found that self management support led to reduced healthcare resource use without any significant change in self management behaviours.²¹⁵ Similar findings are apparent in research about asthma.²¹⁶

Other evidence suggests limited impacts on service use or costs.^{217,218,219,220,221} A systematic review of nine trials of self management education versus usual care for people with chronic obstructive pulmonary disease found that self management education reduced the need for rescue medication and increased courses of oral steroids and antibiotics for respiratory symptoms, but had no effect on hospital admissions, emergency department visits, days lost from work, or lung function.²²²

Similarly, an evaluation of self management courses run in the UK by volunteer tutors at Depression Alliance, Diabetes UK, National Endometriosis Society, Haemophilia Society, British Liver Trust, ME Association, Action for ME, National Osteoporosis Society, British Polio Fellowship, Breakthrough Deaf Integration, and the LINK centre for Deafened People found small improvements in self efficacy, but no significant impacts on the use of health services. The evaluators suggested that disease specific information may need to be included within the generic course structure.²²³

Box 6: contradictory findings

A number of studies have found no or limited effects of supporting self management on healthcare resource use or costs. For instance, a UK cost effectiveness analysis with more than 800 people with arthritis found that a six session self management education programme may not save costs overall or be any more effective than usual care or providing written information alone.²²⁴

Another randomised trial in the UK found that a self management cd rom for children with asthma improved self management, self efficacy and feelings of control but there were no significant reductions in crisis medication use or days off school.²²⁵

A Cochrane review found that education by lay educators may lead to short term improvements in self management attitudes and behaviours but no improvements in healthcare resource use.²²⁶

A randomised trial found that sending reminders to visit GPs improved primary care visits and reduced emergency department visits in 174 children with asthma in Australia.²²⁷ However another trial in the US found that teaching nurses about self management as well as providing patient education did not affect hospitalisations.²²⁸ This demonstrates that the link between supporting self management and outcomes is **complex** and does not necessarily follow a linear pattern.

It also shows that most of the available evidence focuses on self management education courses rather than the wider range of support initiatives available.

However, assuming that supporting self management will reduce healthcare resource use is somewhat simplistic. The aim of supporting self management may be to alter the **pattern of behaviour and service use** rather than reducing service use per se. Therefore it may be more useful to focus on whether different types of services are being used (such as primary care versus secondary care or telephone and online resources versus in person resources).

Overall though, despite some conflicting findings, the evidence suggests that proactively supporting self management and focusing on self efficacy and behaviour change can have an impact on clinical outcomes, crises and unplanned admissions or other costly emergency service use.


WHAT TYPE OF SUPPORT WORKS?

A multitude of approaches have been tested to support self management. These can be conceptualised on a continuum with more passive information sharing approaches at one end and more active behavioural change interventions at the other. Another way to conceptualise self management support is to divide interventions into those that focus on building knowledge and technical skills (such as insulin management) versus those that aim to build self efficacy (confidence in self care). Figure 1 illustrates these typologies and positions various types of self management support along the continuum.

It is difficult to categorise interventions in this way because there is wide variation within interventions. For instance, one type of group education may focus solely on information provision whereas another group education programme may seek to build confidence and change behaviours.

While the placement of each individual intervention is illustrative only, and a matter for debate, conceptualising support approaches as a continuum is useful. Evidence from hundreds of studies suggests that the proactive interventions located in the top right hand quadrant of Figure 1 may be associated with greater change or more sustained levels of behavioural and clinical benefits. Examples of research studies are described in more detail in Appendix 2. This section provides a summary of key concepts only.

Figure 1: continuum of strategies to support self management



Providing information

Providing information about people's condition and how to manage it is an important component of supporting self management. Information can be provided using leaflets, websites, email, text messages, electronic forums, by telephone and in person individually or in groups. Researchers have examined the value of different approaches for supporting people with long term conditions generally as well as subgroups such as children and young people, older people and those from minority ethnic groups and deprived neighbourhoods.

Evidence suggests that providing information verbally or in writing can improve people's knowledge about their condition and care. This may or may not translate into feeling more confident about looking after themselves and improved self management behaviours.^{229,230}

Systematic reviews, randomised trials and other high quality research has found that when used alone, information provision can improve some health behaviours but this does not tend to have lasting follow on effects on clinical outcomes or health service use. However, when used as part of a broader support initiative, information provision has been found to be useful, especially if it is targeted or personalised to account for people's individual needs.²³¹

Box 7: example of tailored information

A systematic review of 60 randomised control trials examined the effectiveness of self help interventions for smoking cessation. The review found that the personalised written materials were more effective than individualised advice from a clinician, which was in turn more effective than non personalised written materials. Adding additional face to face or other advice to tailored written materials did not improve effectiveness compared to personalised materials alone.²³²

Providing support

Another type of self management support involves helping people to change their attitudes, perhaps through group or individual education sessions, care plans, decision support tools or other support mechanisms. This is different from information provision alone because the interventions aim to provide incentives for change or help people learn new skills or practical strategies for coping.

For example, **decision support tools** have been used to support self management.²³³ Such tools may encourage service users and their carers to take more responsibility for their care, help people with long term conditions feel more in control, encourage health professionals to follow recommended care protocols, and have some impacts on quality of life.^{234,235} But reviews about written decision aids suggest that such aids generally affect attitudes and knowledge rather than behaviours.^{236,237}

Care plans and goal setting have also been investigated. A **care plan** is a written document collaboratively designed by service users and professionals covering issues, interventions and review processes.²³⁸ Research suggests that planning care and setting goals can help motivate behaviour change and reduce health service use, but only when there is active and regular follow up.

Courses which provide practical and proactive support have also been found to be more effective for changing people's behaviours and improving clinical outcomes.²³⁹

Furthermore, there is an increasing focus on supporting self care in a way that involves people's carers and family members. Evidence to date tends to focus on children, young people and older people, but there is scope for more detailed examination of engaging families to support self care for a wider range of people.^{240,241}

Behaviour change approaches

The most promising way of supporting self management appears to involve approaches which empower and activate people so they feel more confident about managing their conditions and are more likely to alter their behaviours. There is strong evidence suggesting that improved self efficacy is associated with better clinical outcomes.^{242,243,244}

These approaches include motivational interviewing by telephone or in person, group or individual education programmes with an active component, proactive goal setting and follow up and programmes based on psychological and emotional support which acknowledge people's stage of change.

Box 8: example of stage of change approach

Researchers in the US tested assessing people's capabilities for self management and then tailoring telephone coaching support based on this assessment. Compared to usual disease management, the group receiving tailored support based on their 'stage of change' felt more empowered, had improved clinical outcomes and reduced their use of health services.²⁴⁵

Individual and group education sessions are the most commonly evaluated interventions of this type, though there is also an increasing focus on telephone coaching by nurses.

Randomised trials have found that one off **individual education** visits by nurses or other personnel can be motivating in the short term,²⁴⁶ but that sustained approaches may be more beneficial.^{247,248,249,250,251}

Research suggests that **group education** programmes can improve how people feel about their condition and their ability to cope on a day to day basis.^{252,253,254,255,256,257}

There is also evidence about the potential to improve symptoms,^{258,259,260,261,262,263,264} perceived quality of care,²⁶⁵ and healthcare use,^{266,267,268,269,270,271, 272,273,274,275,276} although findings do vary according to the context, condition, participant characteristics and the extent to which the courses encourage proactive behaviour change.

Self monitoring occurs when people monitor their symptoms or vital signs, perhaps electronically. This allows people to track their progress, modify their behaviours or medications accordingly, and decide when to seek help from health professionals.²⁷⁷ There is evidence that self monitoring can influence behaviour change and have some positive effects on clinical outcomes, but a systematic review found that less than half of people with some long term conditions may be eligible or find this acceptable.²⁷⁸

To summarise, the evidence suggests that not all mechanisms to support self management have equal outcomes. Whilst information provision and building technical skills is important, this is just one facet of self management support. Approaches which recognise people's motivations and needs, take account of their level of desire to change (or stage of change) and support people emotionally and psychologically have been found to have more sustainable impacts on behaviour, clinical outcomes and healthcare resource use.

Further details are available in Appendix 2.

WHAT ELSE DO WE NEED TO KNOW?

Changing behaviours

The evidence is clear that supporting self management can have real benefits for people using services and their families, and the wider health economy. However this is a relatively new area of investigation and knowledge continues to develop.

Interventions to support self management vary considerably in their aims, approach, content, delivery, duration and target group. Even so, there are some general principles that have been found to work well to support self management, including involving people in decision making, proactive education, setting goals and following up on the extent to which these are achieved over time, and helping people manage the social, emotional and physical impacts of their conditions.²⁷⁹

Whilst the general principles are clear, the best **strategies** to achieve this and help motivate people to change their behaviours remain uncertain. Although there is work about smoking cessation, sports psychology and the stages of change model which seeks to examine how to improve self efficacy and motivation to change, the extent to which this applicable to people with long term conditions in the UK is unknown.

A number of innovative strategies are being tested to support behaviour change in the UK, including the Health Foundation's Co-creating Health initiatives. Rigorous evaluation of these programmes and wide dissemination of learning will enhance knowledge in this area considerably.

Supporting clinicians

The **attitudes and skills of healthcare providers** can have a significant effect on the extent to which people feel engaged and supported and this is an area in need of further exploration.^{280,281,282,283,284,285,286}

If people are to be more involved in decisions about their care and more active in keeping themselves well, clinicians need to be able to communicate information effectively and to consider what level of involvement is appropriate for different people.^{287,288} Yet, there may be considerable work to do in this area.^{289,290,291} Interviews with GPs in 11 European countries found that most GPs thought that involving people in healthcare decisions had positive outcomes but GPs defined patient involvement narrowly, thought it took place solely during consultations and felt that they had limited time to engage with people.²⁹² Thus there may be work to do to educate clinicians about the value and scope of supporting self management and the skills they need to achieve this.

Research in the UK suggests that clinicians may take a 'compliance orientated' approach to self management and this is unlikely to be helpful.²⁹³

A number of strategies have been tested to improve clinician communication strategies and help professionals support self management.^{294,295,296,297,298,299,300,301} For instance, a randomised trial found that training GPs about risk communication tools and shared decision making for people with long term conditions could improve prescribing and was unlikely to have major impacts on the cost of care in the UK.³⁰² However another UK study found that while GPs appear receptive to patient involvement, training in shared decision making and risk communication did not help them achieve this or improve patient outcomes.³⁰³ This shows that knowledge is limited about the best strategies to help clinicians support self management.

SUMMARY

Self management works

This review of more than 550 pieces of high quality research suggests that it is worthwhile to support self management, in particular through focusing on behaviour change and supporting self efficacy.

Hundreds of systematic reviews, randomised controlled trials and large observational studies have examined the impact of supporting self management for people with long term conditions. Whilst the findings of individual studies are mixed, the totality of evidence suggests that supporting self management can have benefits for people's attitudes and behaviours, quality of life, clinical symptoms and use of healthcare resources.

Some suggest that the evidence for supporting self management is only moderate but this is because a wide range of things are described as 'self management support' and some may be more effective than others. Past reviews have tended to combine initiatives that focus solely on information provision with interventions that more actively target behaviour change and self efficacy. However, these varying interventions may have different outcomes so combining them could dilute the findings.

Supporting self management has the potential to alleviate the pressure on health and social services caused by workforce shortages, rising demand for services, population increases and budgetary constraints. However, implementing one off interventions is unlikely to make a significant impact on the overall health of the population or on the sustainability of health and social care systems. Supporting self management is not a panacea, and is likely to work best when implemented as part of wider initiatives to improve care through educating practitioners, applying best evidence, and using technology, decision aids and community partnerships effectively.

Proactive strategies work best

There are a wide range of initiatives to support self management. These can be categorised along a continuum of interventions, with passive information provision about healthy behaviours and other 'technical' topics at one end of the scale and initiatives that more actively seek to support behaviour change and increase self efficacy at the other end of the continuum.

Different clinical conditions may require varying approaches to support self management. For instance, people with conditions such as diabetes may benefit from structured education about how to eat, exercise and take medications. For conditions such as depression or chronic pain on the other hand, less 'technical' or clinical education may be needed because the service user has less 'technical work' to do.³⁰⁴ Therefore, evidence about self management support for these groups tends to focus on cognitive and behavioural interventions. Such interventions may be equally valuable for people with diabetes and other conditions, even though the focus to date has been mainly 'technical'.^{305,306,307}

All of the different types of support are important components of the jigsaw needed to encourage self management, but information provision alone is unlikely to be sufficient to motivate sustainable behaviour change and improve clinical outcomes. More active goal setting and behavioural change interventions are necessary. There is also emerging evidence that strategies co-created by service users and professionals or co-led by service users have positive outcomes.^{308,309,310}

"A co-created teaching approach better meets the learning needs of adults with type 2 diabetes mellitus and results in enhanced ability to perform the self care activities required for successful diabetes control."³¹¹

Evidence is developing

Knowledge in this area is developing so evidence about the best strategies to support behaviour change may be limited at this stage,^{312,313} though much work suggests that in order to change behaviour, people need to really want to change.³¹⁴

General components that have been found to work well to support self management include^{315,316,317,318,319,320,321,322,323,324,325,326,327,328,329,330}

- **promoting** healthy lifestyles and **educating** people about their conditions and how to self manage
- **involving** people in decision making
- **motivating** people to self manage using targeted approaches and structured information and support
- helping people **monitor** their symptoms and know when to take appropriate action
- emphasising **problem solving**
- developing care plans as a **partnership** between service users and professionals
- setting **goals** and following up on the extent to which these are achieved over time
- helping people manage the social, emotional and physical **impacts** of their conditions
- proactive **follow up**
- providing opportunities to share and learn from other **service users**

However, the best strategies for implementing these principles and the barriers and facilitators remain uncertain.³³¹ There is a need for high quality research and evaluations that focus on building relationships between service users and practitioners and exploring the most effective strategies for encouraging behaviour change.³³²

Another core component of supporting self management is enabling and encouraging clinicians and lay trainers to work with people to improve their motivation to change.³³³ It is important to understand the skills that clinicians need to help people make changes and the barriers that may stop them from offering such support.^{334,335,336}

Whilst evidence is emerging, there is still a long way to go before we understand the education and support necessary to optimise clinicians' attitudes, skills and behaviours towards self management.³³⁷ This also calls for a fundamental shift in power dynamics and the way both patients and professionals view their roles.^{338,339,340}

“Self management support is the assistance caregivers give patients with chronic disease in order to encourage daily decisions that improve health related behaviors and clinical outcomes. Self management support can be viewed in two ways: as a portfolio of techniques and tools that help patients choose healthy behaviours; and a fundamental transformation of the patient-caregiver relationship into a collaborative partnership.”³⁴¹

Some suggest that training strategies need to account for practitioners' stage of change as well as that of patients.³⁴²

In order to do this we need to understand more about how best to increase self efficacy and encourage behaviour change, and how we can motivate all members of the team, including clinicians and patients themselves to be part of this.

APPENDIX 1 – REVIEW METHODS

To collate evidence, two reviewers searched bibliographic databases, reference lists of identified articles and reviews, and the websites of relevant agencies for information available as at **September 2010**. The search, analysis and narrative synthesis were completed over a three week period. The databases included Medline, Embase, ERIC, Science Citation Index, Cochrane Database of Systematic Reviews, Cochrane Controlled Trials Register, DARE, NHS Health Technology Assessment and Economic Assessment databases, NHS Research Register, NHS Evidence, US National Electronic Library for Health, PsychLit, the WHO Library, Agency for Healthcare Research and Quality, Web of Knowledge, and Ovid.

All databases were searched from inception until present using terms such as self management, self care, self efficacy, self help, self treatment, self monitoring, home monitoring, self medication, support, social support, peer support, mutual support, self management education, long term conditions, chronic care, coping skills, behaviour change, care plans, patient held records, home care, telemedicine and telecare.

To be **eligible** for inclusion in the review, studies had to:

- be primary **research** or reviews
- be **published** research
- be **readily available** online, in print or from relevant organisations
- be available in abstract, journal article, or full report form

Studies in any language were eligible. Randomised controlled trials and systematic reviews were prioritised, though less rigorous designs were also included if few randomised trials or systematic reviews were available.

We scanned more than 100,000 pieces of research, selecting the highest quality and most relevant to summarise here. No formal quality weighting was undertaken.

More than 550 of the highest quality studies and descriptive overviews were synthesised. Data were extracted from all publications using a structured template and studies were grouped according to topic areas and outcomes to provide a **narrative summary** of key trends. Meta analysis was not appropriate given the heterogeneity of the material.

When interpreting the findings it is important to bear in mind several caveats. Firstly, interventions to support self management **vary considerably** in their aims, approach, content, delivery, duration and target group.^{343,344,345} Therefore it would be misleading to refer to 'self management initiatives' as an integrated whole. In the main, this review does not differentiate between various types of interventions because the aim is to examine effects of self management support in all its facets. However, it should not be assumed that all types of self management support have the same impacts or findings.

In most cases there is **limited detail** within research reports about how programmes are implemented. This means that it is difficult to differentiate the most effective components or strategies. Similarly, most available research does not assess the mechanisms by which supporting self management may work. The focus tends to be on outcomes such as quality of life, functional status, clinical outcomes and healthcare resource use. The relationship between a specific self management intervention and these outcomes is explored, but the mechanisms by which these outcomes may occur tend to receive less attention.

A **lack of evidence** or comparisons does not necessarily mean that there is no relationship or benefit, just that there is currently insufficient research to draw conclusions.

Finally, much of the available evidence is sourced from countries with very different healthcare economies and styles of working than the UK so may not be directly **comparable**.

APPENDIX 2 – RESEARCH DETAILS

This appendix describes selected studies about supporting self management in more detail. The aim is to provide a flavour of the scope and nature of the available research.

Evidence suggests that proactive strategies which seek to improve self efficacy and encourage behaviour change are more effective than support which focuses merely on providing information about ‘technical’ issues such as the long term condition or specific self care actions.

Here we provide examples of studies of more passive information sharing through written, computer, video and group education means before moving on to more proactive approaches such as goal setting and telephone coaching.

Written information

Accessible information is a cornerstone of improved self management. A great deal has been written about different ways to provide information to people with long term conditions. We identified more than 60 systematic reviews and randomised trials about providing accessible information through written materials, educational sessions, and technologies such as the internet and video.

A number of written information materials to support self management have been evaluated, including decision aids, guidebooks, and printed educational materials.³⁴⁶ There is some evidence that written **motivational leaflets** or letters can help people feel more confident to raise their concerns and discuss their symptoms,³⁴⁷ but there is sparse evidence that such methods improve self management behaviours or clinical outcomes.^{348,349}

Other reviews suggest that printed materials can improve **knowledge**,^{350,351,352} but may not impact behaviour when used alone.^{353,354}

But findings are mixed. Some trials suggest that postal educational materials are as effective for improving symptoms and self efficacy as group education sessions.³⁵⁵ There is also evidence that combining written information with lectures or other educational activities can be more effective than written information alone.^{356,357}

To understand these differences, it is worth considering the characteristics of the most effective written information tools. There is some evidence that **targeting and personalising** written information is more effective than standardised printed materials.^{358,359,360,361} For example, a randomised trial in Scotland compared posting four personalised asthma education booklets versus conventional oral education at outpatient or surgery visits. Personalised booklets improved self management and reduced hospital admissions.³⁶² Other studies have reinforced these findings.³⁶³

Electronic information sources

Self management support can also be delivered using audiovisual technology, computers, and the mass media.^{364,365,366,367}

There is evidence that providing structured education programmes by **video/dvd, audio or computer** may be as effective as in person education groups.^{368,369} For instance, a randomised trial with older people with long term conditions compared group education, a home study intervention using videotapes and booklets, and a control group. Compared with controls, both educational interventions were associated with reduced pain, sleep difficulties, and symptoms of depression and anxiety. The video course was also associated with reduced symptoms. The authors concluded that lower cost, more accessible home study education using video or dvd may be an effective alternative to group instruction for people with long term conditions.³⁷⁰ Other research has similar findings.³⁷¹

Other novel approaches have been tried. A randomised trial in the US tested an asthma education **video game** as part of a self management programme for high risk children with asthma. The video game was associated with improved quality of life and asthma knowledge.³⁷²

Researchers in many parts of the world have tested using computers to provide self management information and education.^{373,374,375,376,377,378,379,380,381,382,383} For instance, a systematic review of 22 randomised trials found that computerised patient education improved health status in people with various conditions.³⁸⁴ Studies also suggest some benefits for children and young people^{385,386} and less advantaged socio-economic and ethnic groups.³⁸⁷

However while there are some positive trends, other studies have found limited or mixed benefits from online information and support programmes.^{388,389,390}

A number of computer based peer to peer communities and **electronic groups** have been set up to support self management but their effects remain uncertain.³⁹¹ Some descriptive studies suggest that computer chat rooms, coaching and other online forums can provide a good motivator for self care,^{392,393,394} but the effect on clinical outcomes is uncertain.

Another novel approach is using **text messages** or pager messages as reminders and support mechanisms.^{395,396} One trial found that, when combined with other strategies, sending text messages to young people with diabetes helped to improve self management behaviours.³⁹⁷

One randomised trial assessed sending standardised health promotion **email** messages to people weekly for 12 weeks. People who received emails had better self management behaviours and reported favourable changes in healthy eating and physical activity.³⁹⁸ Other studies suggest that email interventions can be targeted to the person's 'stage of change', thus providing more tailored and meaningful support.³⁹⁹

Individual education sessions

A number of studies have assessed the impacts of one to one self management support sessions for people with long term conditions.^{400,401} For instance, a randomised trial in the UK assessed whether specialist asthma nurses could increase knowledge and improve self management during one to one sessions in hospital. People receiving individual education had increased knowledge and less emergency GP call outs in the four months after hospital discharge. Hospital readmission rates were similar between groups.⁴⁰²

Generally, studies suggest that while individual education may increase people's knowledge, it is unlikely to have significant impacts on behaviour and clinical outcomes unless it is targeted, specific, and long term.⁴⁰³

Individualised coaching may be especially beneficial when combined with personalised written materials or other mechanisms to support self management.⁴⁰⁴ For example, a large meta analysis of interventions for people with high blood pressure assessed individual education, self blood pressure monitoring, and structured courses. Individual education was the most effective single strategy for improving blood pressure control, but combining individual education and group sessions was even more effective.⁴⁰⁵ Other reviews have found similarly positive results for people with heart failure,⁴⁰⁶ arthritis,⁴⁰⁷ diabetes,⁴⁰⁸ and asthma.⁴⁰⁹

There has also been success when using individual person centred discussions initially, followed by referral to other self management support strategies (such as educational groups) once needs have been established.⁴¹⁰

Though most individualised counselling has been trialled with nurses or family doctors, pharmacists are increasingly being acknowledged as a valuable resource in supporting self management.⁴¹¹

Group education courses

A great deal has been written about self management group education programmes, including those adapted to specific cultures, demographic groups, learning styles or disease types.^{412,413,414,415,416,417,418,419,420,421,422,423,424,425,426,427}

Group education and peer support programmes aim to help people learn how to manage their own care more effectively, including when to use different healthcare services and resources.^{428,429} More than 60 systematic reviews and numerous additional randomised trials are available in this area.^{430,431} Most take place in health settings or the community, but workplace and family support programmes and school education have also been trialled.^{432,433,434,}

Research generally suggests that group education can improve people's self confidence, clinical outcomes and even health service use. However, not all programmes have been successful, especially with regard to improving symptoms or reducing healthcare use.^{435,436,437,438,439,440,441,442,443}

Reviews and meta analyses are less likely than individual trials to demonstrate a difference but this may be because reviews tend to combine many different interventions and may evaluate interventions using different outcomes than they were set up to achieve.^{444,445}

Education sessions range from those focused on 'technical' information such as how to monitor blood pressure and what foods to eat, to more 'proactive' courses which seek to change people's attitudes towards self management and motivate behaviour change. There is evidence that courses which focus on enhancing self efficacy or combine technical education with more proactive motivation have most effect on clinical outcomes.

Patient held records

A number of strategies have been trialled to increase people's involvement in healthcare processes and decision making as a way of facilitating self management. The main strategies that have been researched include patient held records, care plans and decision support tools.

Sometimes people are given their medical records to keep and bring to each consultation, which is known as **patient held records**. A number of reviews and trials suggest that patient held records have limited effects on self management. For instance, a trial of patient held records for people who had suffered stroke in the US found that while participants were pleased to have a copy of their records, took them when they visited doctors, and reported learning more about their stroke, there was no difference in health practices or behaviours compared to usual care.⁴⁴⁶ Randomised trials in the UK have drawn similar conclusions about patient held records.^{447,448}

There is also interest in making records available electronically for service users.^{449,450,451} A randomised trial in the US provided **patient records online** to people with heart failure. After one year, those who had access to their records online were more likely to adhere to treatment, but there were no differences in self efficacy or satisfaction with care.⁴⁵² This suggests that patient held records may have some impacts on self management strategies, but these impacts are not clear cut. The evidence is too mixed to suggest that patient held records are a useful enabler for self management.

There are similar findings regarding decision support tools.⁴⁵³

Planning and agenda setting

Care plans may include both goal setting and developing plans for how to achieve these goals.⁴⁵⁴ Most of the available evidence about self management treatment plans focuses on people with asthma or COPD, though there is emerging evidence about other conditions too.⁴⁵⁵

There is evidence that action plans or proactive planning support may improve **self management** behaviours for people with long term conditions,^{456,457,458,459,460,461, 462,463,464,465} and this may impact on healthcare resource use.^{466,467,468} Systematic reviews support an agenda setting action plan coupled with self adjustment of medications and regular medical review for people with asthma.^{469,470}

Numerous examples are available. For instance, a randomised trial of 140 adults with asthma or chronic obstructive pulmonary disease assessed developing a written self management plan in groups coupled with individual educational sessions. The plans were associated with improved medication compliance among people with asthma, but not chronic obstructive pulmonary disease.⁴⁷¹

In New Zealand, a randomised trial found similar benefits from written self management plans for children with asthma.⁴⁷²

However, there are questions about whether agenda setting and care planning are supported by practitioners and service users⁴⁷³ or directly impact on clinical outcomes.⁴⁷⁴ A Cochrane review with seven randomised trials examined whether a written asthma self management plan increased adherence to medications and improved clinical outcomes. There was no strong evidence that written plans improved patient outcomes. One type of agenda setting was not consistently more effective than another.⁴⁷⁵

Another review found that adding written self treatment guidelines to self management programmes may improve health outcomes, but the only two controlled studies on this topic found no effect.⁴⁷⁶

A randomised trial in France found that when agenda setting and plans were used as part of a self management programme, only those who adhered fully to the self management plan had improved symptoms.⁴⁷⁷

Evidence about the impact on healthcare resource use is mixed.⁴⁷⁸ A Cochrane review of 36 randomised trials of asthma self management programmes found that combining self monitoring and written agenda setting reduced hospitalisations, emergency department visits, unscheduled visits to the doctor and days off work or school. The reviewers concluded that programmes that enable people to adjust their medication using a written action plan appear to be more effective than other forms of asthma self management.⁴⁷⁹

Stepwise agenda setting and planning has also been found to be beneficial for people with COPD. A randomised trial in Norway found that GP visits reduced by 85% and there was less need for medication and reduced overall health costs during a 12 month follow up period.⁴⁸⁰ Other forms of agenda setting and care planning have also been found to have an impact on healthcare resource use.⁴⁸¹

However other reviews and trials have found limited benefits from agenda setting and care planning,^{482, 483} particularly for those who have been hospitalised.^{484,485,486}

Studies have attempted to explore why action plans and agenda setting seems to work well sometimes and not other times, but few firm conclusions are possible. Plans and agenda setting seem to be better when care plans are provided and supported in primary care compared to secondary care. This approach may be better as a 'preventive' measure rather for those with the most severe disease, or for those who are hospitalised for the first time.⁴⁸⁷

Goal setting and follow up

Self management education often includes some form of goal setting.^{488,489,490,491} A number of studies have outlined the benefits of goal setting as part of self management support.^{492,493,494,495,496,497,498,499,500,501,502}

For instance, a trial in the US found that personalised goal setting as part of a self management support intervention for older women with heart conditions was associated with reduced days in hospital and reduced overall healthcare costs.⁵⁰³

Research has examined the benefits of following up service users to support them in achieving goals and improving self management behaviours.^{504,505,506,507,508,509} For example, a review of workplace self management interventions found that offering opportunities to practice skills and have ongoing follow up were key success factors.⁵¹⁰

A small randomised trial found that self management counselling with goal follow up improved the use of community resources, physical activity and adherence to medication.⁵¹¹ Other trials have found improved self care knowledge and behaviours^{512,513} and reduced hospital admissions and days in hospital when proactive follow up is used as part of self management support.⁵¹⁴

Another trial in the US assessed the cost effectiveness of a brief dietary intervention for people with diabetes. The intervention included touch screen computer assisted assessment, goal setting, and follow up calls and videotape intervention at regular intervals. The intervention was low cost and improved self care behaviour significantly.⁵¹⁵

A trial in the Netherlands examined an electronic diary and follow up device based on the principles of motivational interviewing. When people took part in healthy behaviours, they were positively reinforced by the device. This type of follow up resulted in improved self management.⁵¹⁶

Not all findings about goal setting are positive. As part of a skill orientated self management programme in the Netherlands, people with COPD recorded their symptoms in diaries and graded their health status from 1-10 in a weekly report. A randomised trial found no effect on quality of life or functional status.⁵¹⁷

The thing that seems to make a difference is regular and proactive follow up. In the US, a randomised trial found that the more follow ups and support sessions involved, the more likely people were to control their diabetes.⁵¹⁸ However, another US trial tested whether telephone follow up every month or every three months following self management education might be most effective. There were no significant differences between these follow up intervals on any clinical or quality of life outcomes.⁵¹⁹ This demonstrates that there is still a lot to learn about the best strategies for encouraging behaviour change.

Telephone coaching

Various types of telephone based support and coaching have been tested to encourage self management.^{520,521,522,523}

A number of studies suggest that proactive nurse led telephone calls can be used to encourage self monitoring and self management.^{524,525,526,527,528} For example, a randomised trial with 'high risk' people with diabetes found that automated educational calls with telephone follow up by nurses may improve self care behaviour, glycaemic control, and symptoms among vulnerable people.⁵²⁹ Similar evidence is available from many parts of the world.⁵³⁰

A study in the US found that motivational interviewing helped improve self efficacy, patient activation, lifestyle change and perceived health status.⁵³¹ There are many similar studies about the benefits of motivational interviewing, some of which combine both face to face and telephone approaches.⁵³²

There is also evidence about the benefits of more detailed case management via telephone.⁵³³ A randomised trial in the US evaluated six months of standardised telephone case management for 358 people with heart failure. Telephone case management helped to motivate people to self care, and was associated with about half as many hospitalisations for heart failure, fewer days in hospital, and lower inpatient costs after six months.⁵³⁴

There is evidence to suggest that telephone support is often just as effective as face to face supported self management.^{535,536}

But not all studies are positive. For instance a trial with more than 300 people with diabetes, predominantly from minority ethnic groups, found that clinic based telephonic disease management support in between primary care visits did not improve clinical or behavioural outcomes at one year compared to people receiving usual care.⁵³⁷ Feedback from service providers also suggests there can be practical and technical difficulties.⁵³⁸

Work based support

Work based programmes are gaining increasing recognition for supporting self management, especially in the US where health insurance is often part of employment packages. One of the key differences between workplace programmes and those run by governments or health services is that programmes in the workplace can specifically target a 'captive audience' and may be able to focus on people at higher risk more easily.⁵³⁹

A number of group education and individualised counselling approaches have been tried, whereby employees are followed up regularly in the workplace.^{540,541,542,543,544} Other workplace initiatives include distributing written materials,⁵⁴⁵ providing health risk assessment,⁵⁴⁶ health fairs, weight loss and nutrition classes, fitness programmes, social networks,⁵⁴⁷ online support⁵⁴⁸ and environmental changes such as introducing walking tracks around the work environment.⁵⁴⁹

While there is little evidence about the most effective workplace strategy, most research suggests that these initiatives can help to support self management behaviours. One review of 35 different interventions found that offering opportunities to practice skills and have ongoing follow up were more effective than education alone.⁵⁵⁰ Another review found that providing individualised risk reduction programmes for high risk employees is a critical element of successful worksite interventions.⁵⁵¹ This emphasises that regardless of the setting, proactive interventions appear more effective for encouraging self management behaviours and resulting changes in clinical outcomes.

Self monitoring and telemonitoring

Self monitoring involves service users monitoring their symptoms in order to track their progress, modify their behaviours or medications accordingly, or assess when to seek help from health professionals.⁵⁵² Self monitoring is often linked with electronic monitoring devices, but this term can also refer to written management plans and systems to help patients self refer to health services.⁵⁵³ We identified more than 50 studies about self monitoring, most of which had positive outcomes, though not all were of high quality.^{554,555,556,557,558,559,560}

Randomised trials suggest that electronic self monitoring may have some clinical benefits.^{561,562,563} For instance, a trial in Germany found that a self monitoring blood glucose device for people with diabetes improved glycaemic control and general wellbeing.⁵⁶⁴ Other trials support the value of self monitoring for people with diabetes,^{565,566,567} asthma,^{568,569,570} and high blood pressure^{571,572} amongst others.^{573,574,575}

However, there are some conflicting findings.⁵⁷⁶ A trial in Scotland concluded that prescribing peak flow meters and giving self management guidelines to everyone with asthma is unlikely to improve mortality or morbidity.⁵⁷⁷ Another UK trial found that self monitoring did not improve self management or symptoms in children with asthma.⁵⁷⁸ It appears that those most likely to benefit from self monitoring have the most severe disease.

Although there is evidence that self monitoring can have some impacts on clinical outcomes, there is limited evidence about the effect on resource use.⁵⁷⁹

Self monitoring may also be inappropriate or unfeasible some. A Cochrane review found that compared to standard monitoring, people who self monitor can improve the quality of their oral anticoagulation therapy and resulting clinical outcomes. However, self monitoring or self management were not feasible for up to half of the people requiring anticoagulant therapy due to patient refusal, exclusion by their GP, or inability to complete training.⁵⁸⁰

Schemes which use telecommunications systems such as the internet or telephone lines to transfer or record monitoring information are often referred to as 'telemonitoring.' This is not always strictly 'self monitoring' as it may involve interaction between service users and health professionals,^{581,582} but it is another way to support self management.

There is mixed evidence about the value of telemonitoring,⁵⁸³ but it is well received by patients and providers.^{584,585} Most high quality information about the effect of telemonitoring on clinical outcomes focuses on people with diabetes or hypertension, although studies with other patient groups are emerging.⁵⁸⁶ Most available evidence suggests that transmitting data about diabetic symptoms or blood pressure via telephone lines can improve self management and have a positive effect on clinical outcomes.^{587,588,589,590,591,592} This is especially true when the system is linked to nurses or doctors who review the information and provide feedback to patients or take action when needed.

REFERENCES

- 1 World Health Organisation (2002). *Innovative Care for Chronic Conditions. Building Blocks for Action*. Geneva: WHO.
- 2 World Health Organisation (2000). 'Prevention and control of noncommunicable diseases.' *Proceedings of the World Health Assembly*, 15-20 May 2000. Geneva: WHO.
- 3 Department of Health (2004). *Improving Chronic Disease Management*. London: Department of Health.
- 4 Department of Health (2004). *Chronic disease management: A compendium of information*. London: Department of Health.
- 5 Department of Health (2004). *Improving Chronic Disease Management*. London: Department of Health.
- 6 Vickery DM et al (1983). 'Effect of a self care education program on medical visits.' *JAMA* 250 (21): 2952-2956.
- 7 Department of Health (2005). *Self care - A Real Choice*. London: Department of Health.
- 8 Heisler M (2010). 'Different models to mobilize peer support to improve diabetes self-management and clinical outcomes: evidence, logistics, evaluation considerations and needs for future research.' *Fam Pract* 27(Suppl 1): i23-32.
- 9 Dennis SM, Zwar N, Griffiths R, et al (2008). 'Chronic disease management in primary care: from evidence to policy.' *Med J Aust* 188(Suppl 8): S53-S56.
- 10 Padilha JM (2010). 'Influence of perception of chronic obstructive pulmonary disease in promotion of self-management of disease.' *Rev Port Pneumol* 16(4): 641-8.
- 11 Lubetkin EI, Lu WH, Gold MR (2010). 'Levels and correlates of patient activation in health center settings: building strategies for improving health outcomes.' *J Health Care Poor Underserved* 21(3): 796-808.
- 12 Shigaki C, Kruse RL, Mehr D, Sheldon KM, Bin Ge, Moore C, Lemaster J (2010). 'Motivation and diabetes self-management.' *Chronic Illn* 6(3): 202-14.
- 13 Kaptein AA, Klok T, Moss-Morris R, Brand PL (2010). 'Illness perceptions: impact on self-management and control in asthma.' *Curr Opin Allergy Clin Immunol* 10(3): 194-9.
- 14 Gallagher R (2010). 'Self management, symptom monitoring and associated factors in people with heart failure living in the community.' *Eur J Cardiovasc Nurs* 9(3): 153-60.
- 15 Coulter A and Ellins J (2006). *Patient-focused Interventions: A review of the Evidence*. London: The Health Foundation.
- 16 Remmers C, Hibbard J, Mosen DM, Wagenfield M, Hoyer RE, Jones C (2009). 'Is patient activation associated with future health outcomes and healthcare utilization among patients with diabetes?' *J Ambul Care Manage* 32(4):320-7.
- 17 Shon KH, Park SS (2002). 'Medication and symptom management education program for the rehabilitation of psychiatric patients in Korea: the effects of promoting schedule on self-efficacy theory.' *Yonsei Med J* 43(5): 579-89.
- 18 Greenfield S, Kaplan S, Ware J E Jr (1985). 'Expanding patient involvement in care. Effects on patient outcomes.' *Ann Intern Med* 102(4): 520-528.
- 19 Blissmer B, Prochaska JO, Velicer WF, Redding CA, Rossi JS, Greene GW, Paiva A, Robbins M (2010). 'Common factors predicting long-term changes in multiple health behaviors.' *J Health Psychol* 15(2): 205-14.
- 20 Song M (2010). 'Diabetes mellitus and the importance of self-care.' *J Cardiovasc Nurs* 25(2): 93-8.
- 21 Fishman L, Barendse RM, Hait E, Burdick C, Arnold J (2010). 'Self-Management of Older Adolescents With Inflammatory Bowel Disease: A Pilot Study of Behavior and Knowledge as Prelude to Transition.' *Clin Pediatr* (published online September 2010).
- 22 Mancuso CA, Sayles W, Allegrante JP (2010). 'Knowledge, Attitude, and Self-Efficacy in Asthma Self-Management and Quality of Life.' *J Asthma* (published online September 2010).
- 23 Sol BG, van der Graaf Y, van Petersen R, Visseren FL (2010). 'The effect of self-efficacy on cardiovascular lifestyle.' *Eur J Cardiovasc Nurs* (published online July 2010).
- 24 Chih AH, Jan CF, Shu SG, Lue BH (2010). 'Self-efficacy affects blood sugar control among adolescents with type I diabetes mellitus.' *J Formos Med Assoc* 109(7): 503-10.
- 25 Du S, Yuan C (2010). 'Evaluation of patient self-management outcomes in health care: a systematic review.' *Int Nurs Rev* 57(2): 159-67.
- 26 Weng LC, Dai YT, Huang HL, Chiang YJ (2010). 'Self-efficacy, self-care behaviours and quality of life of kidney transplant recipients.' *J Adv Nurs* 66(4): 828-38.
- 27 King DK, Glasgow RE, Toobert DJ, Strycker LA, Estabrooks PA, Osuna D, Faber AJ (2010). 'Self-efficacy, problem solving, and social-environmental support are associated with diabetes self-management behaviors.' *Diabetes Care* 33(4): 751-3.
- 28 Cheung C, Wyman J, Gross C, Peters J, Findorff M, Stock H (2010). 'Exercise behavior in older adults: a test of the transtheoretical model.' *J Aging Phys Act* 15(1): 103-18.
- 29 Giloth BE (1990). 'Promoting patient involvement: educational, organizational, and environmental strategies.' *Patient Educ Couns* 15(1): 29-38.
- 30 Hibbard JH, Mahoney ER, Stock R, Tusler M (2007). 'Do increases in patient activation result in improved self management behaviors?' *Health Serv Res* 42(4): 1443-63.
- 31 Rygg LO, Rise MB, Lomundal B, Solberg HS, Steinsbekk A (2010). 'Reasons for participation in group-based type 2 diabetes self-management education. A qualitative study.' *Scand J Public Health* (published online September 2010).
- 32 Department of Health and Diabetes UK (2006). *Care Planning in Diabetes*. Report from the joint Department of Health and Diabetes UK Care Planning Working Group.
- 33 Department of Health (2006). *Supporting People with Long-term Conditions to Self care: A guide to developing local strategies and good practice*. London: Department of Health.
- 34 Yip YB, Sit JW, Fung KK, Wong DY, Chong SY, Chung LH, Ng TP (2007). 'Effects of a self management arthritis programme with an added exercise component for osteoarthritic knee: randomized controlled trial.' *J Adv Nurs* 59(1): 20-8.
- 35 Sturt J, Taylor H, Docherty A, Dale J, Louise T (2006). 'A psychological approach to providing self management education for people with type 2 diabetes: the Diabetes Manual.' *BMC Fam Pract* 7: 70.
- 36 Samoocha D, Bruinvels DJ, Elbers NA, Anema JR, van der Beek AJ (2010). 'Effectiveness of web-based interventions on patient empowerment: a systematic review and meta-analysis.' *J Med Internet Res* 12(2): e23.
- 37 Crawford Shearer NB, Fleury JD, Belyea M (2010). 'Randomized control trial of the Health Empowerment Intervention: feasibility and impact.' *Nurs Res* 59(3): 203-11.

- 38 Klainin P, Ounnapiruk L (2010). 'A meta-analysis of self-care behavior research on elders in Thailand: an update.' *Nurs Sci Q* 23(2): 156-63.
- 39 Greenhalgh T, Collard A, Campbell-Richards D, Vijayaraghavan S, Malik F, Morris J, Claydon A (2010). 'Storylines of self-management: narratives of people with diabetes from a multiethnic inner city population.' *J Health Serv Res Policy* (published online September 2010).
- 40 Jones F, Riazi A (2010). 'Self-efficacy and self-management after stroke: a systematic review.' *Disabil Rehabil* (published online August 2010).
- 41 Wagner JL, Smith G, Ferguson P, van Bakergem K, Hrisko S (2010). 'Pilot study of an integrated cognitive-behavioral and self-management intervention for youth with epilepsy and caregivers: Coping Openly and Personally with Epilepsy (COPE).' *Epilepsy Behav* 18(3): 280-5.
- 42 Barlow JH, Turner, Wright (1998). 'Sharing, caring and learning to take control: self management training for people with arthritis.' *Psy, Health Med* 3(4): 387-93.
- 43 Barlow JH, Turner, Wright (1998). 'Long-term outcomes of an arthritis self management programme.' *Brit J Rheumatology* 37: 1315-9.
- 44 Barlow JH, Turner, Wright (2000). 'A randomised controlled study of the arthritis self management programme in the UK.' *Health Ed Res* 15(6): 665-80.
- 45 Barlow JH, Turner, Wright (2001). 'Patient education for people with arthritis in rural communities: the UK experience.' *Patient Ed Couns* 44: 205-14.
- 46 Lorig KR, Sobel DS, Stewart AL, et al (1999). 'Evidence suggesting that a chronic disease self management program can improve health status while reducing utilization and costs: A randomized trial.' *Med Care* 37(1): 5-14.
- 47 Chiang LC, Huang JL, Yeh KW, Lu CM (2004). 'Effects of a self management asthma educational program in Taiwan based on PRECEDE-PROCEED model for parents with asthmatic children.' *J Asthma* 41(2): 205-15.
- 48 Yu PL, Ye W, Liu XR, et al (2003). 'Evaluation on the effectiveness for self management of hypertensive patients in a community.' *Zhonghua Liu Xing Bing Xue Za Zhi* 24(9): 790-3.
- 49 Bodenheimer T, Lorig K, Holman H, Grumbach K (2002). 'Patient self management of chronic disease in primary care.' *JAMA* 288(19): 2469-75.
- 50 Gadisseur AP, Breukink-Engbers WG, van der Meer FJ, van den Besselaar AM, Sturk A, Rosendaal FR (2003). 'Comparison of the quality of oral anticoagulant therapy through patient self management and management by specialized anticoagulation clinics in the Netherlands: a randomized clinical trial.' *Arch Intern Med* 163(21): 2639-46.
- 51 Gustavsson C, Denison E, Koch L (2010). 'Self-management of persistent neck pain: a randomized controlled trial of a multi-component group intervention in primary health care.' *Eur J Pain* 14(6): 630-e1,e11.
- 52 Parchman ML, Arambula-Solomon TG, Noel PH, Larne AC, Pugh JA (2010). 'Stage of change advancement for diabetes self-management behaviors and glucose control.' *Diabetes Educ* 29(1): 128-34.
- 53 Gorczynski P, Faulkner G, Greening S, Cohn T (2010). 'Exploring the construct validity of the transtheoretical model to structure physical activity interventions for individuals with serious mental illness.' *Psychiatr Rehabil J* 34(1): 61-4.
- 54 Daley LK, Fish AF, Frid DJ, Mitchell GL (2010). 'Stage-specific education/counseling intervention in women with elevated blood pressure.' *Prog Cardiovasc Nurs* 24(2): 45-52.
- 55 Chung SJ, Hoerr SL (2010). 'Evaluation of a theory-based community intervention to increase fruit and vegetable intakes of women with limited incomes.' *Nutr Res Pract* 1(1): 46-51.
- 56 Jackson R, Asimakopoulou K, Scammell A (2010). 'Assessment of the transtheoretical model as used by dietitians in promoting physical activity in people with type 2 diabetes.' *J Hum Nutr Diet* 20(1): 27-36.
- 57 Mochari-Greenberger H, Terry MB, Mosca L (2010). 'Does stage of change modify the effectiveness of an educational intervention to improve diet among family members of hospitalized cardiovascular disease patients?' *J Am Diet Assoc* 110(7): 1027-35.
- 58 Milan JE, White AA (2010). 'Impact of a stage-tailored, web-based intervention on folic Acid-containing multivitamin use by college women.' *Am J Health Promot* 24(6): 388-95.
- 59 Glasson C, Chapman K, James E (2010). 'Fruit and vegetables should be targeted separately in health promotion programmes: differences in consumption levels, barriers, knowledge and stages of readiness for change.' *Public Health Nutr* 25: 1-8.
- 60 Blissmer B, Prochaska JO, Velicer WF, Redding CA, Rossi JS, Greene GW, Paiva A, Robbins M (2010). 'Common factors predicting long-term changes in multiple health behaviors.' *J Health Psychol* 15(2): 205-14.
- 61 Callaghan P, Khalil E, Morres I (2010). 'A prospective evaluation of the Transtheoretical Model of Change applied to exercise in young people.' *Int J Nurs Stud* 47(1): 3-12.
- 62 Hildebrand DA, Betts NM (2010). 'Assessment of stage of change, decisional balance, self-efficacy, and use of processes of change of low-income parents for increasing servings of fruits and vegetables to preschool-aged children.' *J Nutr Educ Behav* 41(2): 110-9.
- 63 Wolf RL, Lepore SJ, Vandergrift JL, Wetmore-Arkader L, McGinty E, Pietrzak G, Yaroch AL (2010). 'Knowledge, barriers, and stage of change as correlates of fruit and vegetable consumption among urban and mostly immigrant black men.' *J Am Diet Assoc* 108(8): 1315-22.
- 64 Greaney ML, Riebe D, Ewing Garber C, Rossi JS, Lees FD, Burbank PA, Nigg CR, Ferrone CL, Clark PG (2010). 'Long-term effects of a stage-based intervention for changing exercise intentions and behavior in older adults.' *Gerontologist* 48(3): 358-67.
- 65 Rau J, Teichmann J, Petermann J (2010). 'Motivation for exercise of cancer patients--a longitudinal study of the Transtheoretical Model (TTM).' *Psychother Psychosom Med Psychol* 58(2): 58-62.
- 66 Helitzer DL, Peterson AB, Sanders M, Thompson J (2010). 'Relationship of stages of change to attendance in a diabetes prevention program.' *Am J Health Promot* 21(6): 517-20.
- 67 Levesque DA, Cummins CO, Prochaska JM, Prochaska JO (2010). 'Randomized trial of stage-based interventions for informed Medicare choices.' *Health Care Financ Rev* 27(4): 25-40.
- 68 Hadjistavropoulos H, Shymkiw J (2010). 'Predicting readiness to self-manage pain.' *Clin J Pain* 23(3): 259-66.
- 69 Tsay SL, Hung LO (2004). 'Empowerment of patients with end-stage renal disease - a randomized controlled trial.' *Int J Nurs Stud* 41(1): 59-65.
- 70 Fu D, Fu H, McGowan P, Shen YE, Zhu L, Yang H, Mao J, Zhu S, Ding Y, Wei Z (2003). 'Implementation and quantitative evaluation of chronic disease self management programme in Shanghai, China: randomized controlled trial.' *Bull World Health Organ* 81(3): 174-82.
- 71 Department of Health (2001). *The Expert Patient: A New Approach to Chronic Disease Management for the 21st Century*. London: Department of Health.
- 72 Schiel R, Braun A, Muller R, et al (2004). 'A structured treatment and educational program for patients with type 2 diabetes mellitus, insulin therapy and impaired cognitive function (DikoL).' *Med Klin* 99(6): 285-92.
- 73 LeFort SM (2000). 'A test of Braden's Self-Help Model in adults with chronic pain.' *J Nursing Scholarship* 32(2): 153-60.

- 74 Damush TM, Weinberger M, Perkins SM, Rao JK, Tierney WM, Qi R, Clark DO (2003). 'The long-term effects of a self management program for inner-city primary care patients with acute low back pain.' *Arch Intern Med* 163(21): 2632-8.
- 75 Polonsky WH, Earles J, Smith S, Pease DJ, Macmillan M, Christensen R, Taylor T, Dickert J, Jackson RA (2003). 'Integrating medical management with diabetes self management training: a randomized control trial of the Diabetes Outpatient Intensive Treatment program.' *Diabetes Care* 26(11): 3048-53.
- 76 Guevara JP, Wolf FM, Grum CM, Clark NM (2003). 'Effects of educational interventions for self management of asthma in children and adolescents: systematic review and meta-analysis.' *BMJ* 326(7402): 1308-9.
- 77 Damush TM, Weinberger M, Perkins SM, Rao JK, Tierney WM, Qi R, Clark DO (2003). 'Randomized trial of a self management program for primary care patients with acute low back pain: short-term effects.' *Arthritis Rheum* 49(2): 179-86.
- 78 Scain SF, Friedman R, Gross JL (2009). 'A structured educational program improves metabolic control in patients with type 2 diabetes: a randomized controlled trial.' *Diabetes Educ* 35(4): 603-11.
- 79 Butz AM, Syron L, Johnson B, Spaulding J, Walker M, Bollinger ME (2005). 'Home-based asthma self management education for inner city children.' *Public Health Nurs* 22(3): 189-99.
- 80 Alp A, Kanat E, Yurtkuran M (2007). 'Efficacy of a self management program for osteoporotic subjects.' *Am J Phys Med Rehabil* 86(8): 633-40.
- 81 Nitzke S, Kritsch K, Boeckner L, Greene G, Hoerr S, Horacek T, Kattelmann K, Lohse B, Oakland MJ, Beatrice P, White A (2010). 'A stage-tailored multi-modal intervention increases fruit and vegetable intakes of low-income young adults.' *Am J Health Promot* 22(1): 6-14.
- 82 Brown SA (1990). 'Studies of educational interventions and outcomes in diabetic adults: a meta-analysis revisited.' *Patient Educ Couns* 16(3): 189-215.
- 83 Barlow JH, Wright CC, Sheasby, et al (2002). 'Self management approaches for people with chronic conditions: a review.' *Patient Ed Couns* 48: 177-87.
- 84 The British Liver Trust (1999). *Living a Healthy Life with Long-term Illness*. Leland Stanford Junior University.
- 85 Millar A, Cauch-Dudek K, Shah BR (2010). 'The impact of diabetes education on blood glucose self-monitoring among older adults.' *J Eval Clin Pract* 16(4): 790-3.
- 86 Tang TS, Funnell MM, Brown MB, Kurlander JE (2010). 'Self-management support in "real-world" settings: an empowerment-based intervention.' *Patient Educ Couns* 79(2): 178-84.
- 87 Bourbeau J, Nault D, Dang-Tan T (2004). 'Self management and behaviour modification in COPD.' *Patient Educ Couns* 52(3): 271-7.
- 88 Yap TL, Brown C, Cromwell DA, van der Meulen J, Emberton M (2009). 'The impact of self management of lower urinary tract symptoms on frequency-volume chart measures.' *BJU Int* 104(8): 1104-8.
- 89 Brody BL, Roch-Levecq AC, Kaplan RM, Moutier CY, Brown SI (2006). 'Age-related macular degeneration: self management and reduction of depressive symptoms in a randomized, controlled study.' *J Am Geriatr Soc* 54(10): 1557-62.
- 90 Glasgow RE, Toobert DJ, Barrera M Jr, Strycker LA (2004). 'Assessment of problem-solving: a key to successful diabetes self management.' *J Behav Med* 27(5): 477-90.
- 91 Wright SP, Walsh H, Ingley KM, et al (2003). 'Uptake of self management strategies in a heart failure management programme.' *Eur J Heart Fail* 5(3): 371-80.
- 92 Kremers IP, Steverink N, Albersnagel FA, Slaets JP (2006). 'Improved self management ability and well-being in older women after a short group intervention.' *Aging Ment Health* 10(5): 476-84.
- 93 Kroenke K, Bair MJ, Damush TM, Wu J, Hoke S, Sutherland J, Tu W (2009). 'Optimized antidepressant therapy and pain self management in primary care patients with depression and musculoskeletal pain: a randomized controlled trial.' *JAMA* 301(20): 2099-2110.
- 94 Krieger J, Takaro TK, Song L, Beaudet N, Edwards K (2009). 'A randomized controlled trial of asthma self management support comparing clinic-based nurses and in-home community health workers: the Seattle-King County Healthy Homes II Project.' *Arch Pediatr Adolesc Med* 163(2): 141-9.
- 95 Barlow JH, et al (2000). *Self management literature review*. Psychosocial Research Centre, Coventry University.
- 96 Cedraschi C, Desmeules J, Rapiti E, Baumgartner E, Cohen P, Finckh A, Allaz AF, Vischer TL (2004). 'Fibromyalgia: a randomised, controlled trial of a treatment programme based on self management.' *Ann Rheum Dis* 63(3): 290-6.
- 97 Wattana C, Srisuphan W, Pothiban L, Upchurch SL (2007). 'Effects of a diabetes self management program on glycemic control, coronary heart disease risk, and quality of life among Thai patients with type 2 diabetes.' *Nurs Health Sci* 9(2): 135-41.
- 98 Smith TR, Nicholson RA, Banks JW (2010). 'Migraine education improves quality of life in a primary care setting.' *Headache* 50(4): 600-12.
- 99 Mancuso CA, Sayles W, Allegrante JP (2010). 'Randomized trial of self-management education in asthmatic patients and effects of depressive symptoms.' *Ann Allergy Asthma Immunol* 105(1): 12-9.
- 100 Didjurgeit U, Kruse J, Schmitz N, Stückenschneider P, Sawicki PT (2002). 'A time-limited, problem-orientated psychotherapeutic intervention in Type 1 diabetic patients with complications: a randomized controlled trial.' *Diabet Med* 19(10): 814-21.
- 101 Furmark T, Carlbring P, Hedman E, Sonnenstein A, Clevberger P, Bohman B, Eriksson A, Hallen A, Frykman M, Holmstrom A, Sparthan E, Tillfors M, Ihrfelt EN, Spak M, Eriksson A, Ekselius L, Andersson G (2009). 'Guided and unguided self-help for social anxiety disorder: randomised controlled trial.' *Br J Psychiatry* 195(5): 440-7.
- 102 Cochran J, Conn VS (2008). 'Meta-analysis of quality of life outcomes following diabetes self management training.' *Diabetes Educ* 34(5): 815-23.
- 103 Korstjens I, May AM, van Weert E, Mesters I, Tan F, Ros WJ, Hoekstra-Weebers JE, van der Schans CP, van den Borne B (2008). 'Quality of life after self management cancer rehabilitation: a randomized controlled trial comparing physical and cognitive-behavioral training versus physical training.' *Psychosom Med* 70(4): 422-9.
- 104 Coleman S, Briffa NK, Carroll G, Inderjeeth C, Cook N, McQuade J (2008). 'Effects of self management, education and specific exercises, delivered by health professionals, in patients with osteoarthritis of the knee.' *BMC Musculoskelet Disord* 9: 133.
- 105 Pramuka M, Hendrickson R, Zinski A, Van Cott AC (2007). 'A psychosocial self management program for epilepsy: a randomized pilot study in adults.' *Epilepsy Behav* 11(4): 533-45.
- 106 Hughes RB, Robinson-Whelen S, Taylor HB, Hall JW (2006). 'Stress self management: an intervention for women with physical disabilities.' *Women's Health Issues* 16(6): 389-99.
- 107 Frieswijk N, Steverink N, Buunk BP, Slaets JP (2006). 'The effectiveness of a bibliotherapy in increasing the self management ability of slightly to moderately frail older people.' *Patient Educ Couns* 61(2): 219-27.

- 108 van Hooff ML, van der Merwe JD, O'Dowd J, Pavlov PW, Spruit M, de Kleuver M, van Limbeek J (2010). 'Daily functioning and self-management in patients with chronic low back pain after an intensive cognitive behavioral programme for pain management.' *Eur Spine J* 19(9): 1517-26.
- 109 Mérelle SY, Sorbi MJ, van Doornen LJ, Passchier J (2008). 'Lay trainers with migraine for a home-based behavioral training: a 6-month follow up study.' *Headache* 48(9): 1311-25.
- 110 McGillion MH, Watt-Watson J, Stevens B, Lefort SM, Coyte P, Graham A (2008). 'Randomized controlled trial of a psychoeducation program for the self management of chronic cardiac pain.' *J Pain Symptom Manage* 36(2): 126-40.
- 111 Devos-Comby L, Cronan T, Roesch SC (2006). 'Do exercise and self management interventions benefit patients with osteoarthritis of the knee? A meta-analytic review.' *J Rheumatol* 33(4): 744-56.
- 112 Cimprich B, Janz NK, Northouse L, Wren PA, Given B, Given CW (2005). 'Taking CHARGE: A self management program for women following breast cancer treatment.' *Psychooncology* 14(9): 704-17.
- 113 Heitkemper MM, Jarrett ME, Levy RL, Cain KC, Burr RL, Feld A, Barney P, Weisman P (2004). 'Self management for women with irritable bowel syndrome.' *Clin Gastroenterol Hepatol* 2(7): 585-96.
- 114 Tang TS, Funnell MM, Brown MB, Kurlander JE (2010). 'Self-management support in "real-world" settings: an empowerment-based intervention.' *Patient Educ Couns* 79(2): 178-84.
- 115 Clark MM, Novotny PJ, Patten CA, Rausch SM, Garces YI, Jatoi A, Sloan JA, Yang P (2010). 'Motivational readiness for physical activity and quality of life in long-term lung cancer survivors.' *Lung Cancer* 61(1): 117-22.
- 116 Yu PL, Ye W, Liu XR, Liu YJ, Zhang J, Bai XL, Wu ZL (2003). 'Evaluation on the effectiveness for self management of hypertensive patients in a community.' *Zhonghua Liu Xing Bing Xue Za Zhi* 24(9): 790-3.
- 117 Siu AM, Chan CC, Poon PK, Chui DY, Chan SC (2007). 'Evaluation of the chronic disease self management program in a Chinese population.' *Patient Educ Couns* 65(1): 42-50.
- 118 Brody BL, Roch-Levecq AC, Gamst AC, Maclean K, Kaplan RM, Brown SI (2002). 'Self management of age-related macular degeneration and quality of life: a randomized controlled trial.' *Arch Ophthalmol* 120(11): 1477-83.
- 119 Lewin RJ, Furze G, Robinson J, Griffith K, Wiseman S, Pye M, Boyle R (2002). 'A randomised controlled trial of a self management plan for patients with newly diagnosed angina.' *Br J Gen Pract* 52(476): 194-6, 199-201.
- 120 Foster G, Taylor SJ, Eldridge SE, Ramsay J, Griffiths CJ (2007). 'Self management education programmes by lay leaders for people with chronic conditions.' *Cochrane Database Syst Rev* (4): CD005108.
- 121 Perneger TV, Sudre P, Muntner P, Uldry C, Courtehouse C, Naef AF, Jacquemet S, Nicod L, Rochat T, Assal JP (2002). 'Effect of patient education on self management skills and health status in patients with asthma: a randomized trial.' *Am J Med* 113(1): 7-14.
- 122 Jones F, Mandy A, Partridge C (2009). 'Changing self-efficacy in individuals following a first time stroke: preliminary study of a novel self management intervention.' *Clin Rehabil* 23(6): 522-33.
- 123 Bernard-Bonnin AC, Stachenko S, Bonin D, et al (1995). 'Self management teaching programs and morbidity of pediatric asthma: a meta-analysis.' *J Allergy Clin Immunol* 95(1 Pt 1): 34-41.
- 124 Wolf FM, Guevara JP, Grum CM, et al (2003). 'Educational interventions for asthma in children.' *Cochrane Database Syst Rev* (1): CD000326.
- 125 Norris SL, Nichols PJ, Caspersen CJ, et al (2002). 'Increasing diabetes self management education in community settings. A systematic review.' *Am J Prev Med* 22(4 Suppl): 39-66.
- 126 Barbanel D, Eldridge S, Griffiths C (2003). 'Can a self management programme delivered by a community pharmacist improve asthma control? A randomised trial.' *Thorax* 58(10): 851-4.
- 127 Foster G, Taylor SJC, Eldridge S, Ramsay J, Griffiths CJ (2007). 'Self management education programmes by lay leaders for people with chronic conditions.' *Cochrane Database of Systematic Reviews* (4).
- 128 Heuts PH, de Bie R, Drieteleaar M, Aretz K, Hopman-Rock M, Bastiaenen CH, Metsemakers JF, van Weel C, van Schayck O (2005). 'Self management in osteoarthritis of hip or knee: a randomized clinical trial in a primary healthcare setting.' *J Rheumatol* 32(3): 543-9.
- 129 Hurley MV, Walsh NE, Mitchell HL, Pimm TJ, Patel A, Williamson E, Jones RH, Dieppe PA, Reeves BC (2007). 'Clinical effectiveness of a rehabilitation program integrating exercise, self management, and active coping strategies for chronic knee pain: a cluster randomized trial.' *Arthritis Rheum* 57(7): 1211-9.
- 130 Brown CS, Wan J, Bachmann G, Rosen R (2009). 'Self management, amitriptyline, and amitriptyline plus tramcinolone in the management of vulvodynia.' *J Womens Health* 18(2): 163-9.
- 131 Sol BG, van der Graaf Y, Brouwer B, Hickox SM, Visseren FL (2010). 'The effect of a self-management intervention to reduce vascular risk factors in patients with manifestations of vascular diseases.' *Eur J Cardiovasc Nurs* 9(2): 132-9.
- 132 Anderson RM, Funnell MM, Butler PM, et al (1995). 'Patient empowerment. Results of a randomized controlled trial.' *Diabetes Care* 18: 943-49.
- 133 Wantland DJ, Holzemer WL, Moezzi S, Willard SS, Arudo J, Kirksey KM, Portillo CJ, Corless IB, Rosa ME, Robinson LL, Nicholas PK, Hamilton MJ, Sefcik EF, Human S, Rivero MM, Maryland M, Huang E (2008). 'A randomized controlled trial testing the efficacy of an HIV/AIDS symptom management manual.' *J Pain Symptom Manage* 36(3): 235-46.
- 134 Yardley L, Kirby S (2006). 'Evaluation of booklet-based self management of symptoms in Ménière disease: a randomized controlled trial.' *Psychosom Med* 68(5): 762-9.
- 135 Davies MJ, Heller S, Skinner TC, Campbell MJ, Carey ME, Cradock S, Dallosso HM, Daly H, Doherty Y, Eaton S, Fox C, Oliver L, Rantell K, Rayman G, Khunti K (2008). 'Effectiveness of the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cluster randomised controlled trial.' *BMJ* 336(7642): 491-5.
- 136 Gregg JA, Callaghan GM, Hayes SC, Glenn-Lawson JL (2007). 'Improving diabetes self management through acceptance, mindfulness, and values: a randomized controlled trial.' *J Consult Clin Psychol* 75(2): 336-43.
- 137 Fitzmaurice DA, Murray ET, McCahon D, Holder R, Raftery JP, Hussain S, Sandhar H, Hobbs FD (2005). 'Self management of oral anticoagulation: randomised trial.' *BMJ* 331(7524): 1057.
- 138 Koertke H, Zittermann A, Wagner O, Koerfer R (2007). 'Self management of oral anticoagulation therapy improves long-term survival in patients with mechanical heart valve replacement.' *Ann Thorac Surg* 83(1): 24-9.
- 139 Goeppinger J, Armstrong B, Schwartz T, Ensley D, Brady TJ (2007). 'Self management education for persons with arthritis: Managing comorbidity and eliminating health disparities.' *Arthritis Rheum* 57(6): 1081-8.
- 140 Warwick M, Gallagher R, Chenoweth L, Stein-Parbury J (2010). 'Self-management and symptom monitoring among older adults with chronic obstructive pulmonary disease.' *J Adv Nurs* 66(4): 784-93.

- 141 Rooks DS, Gautam S, Romeling M, Cross ML, Stratigakis D, Evans B, Goldenberg DL, Iversen MD, Katz JN (2007). 'Group exercise, education, and combination self management in women with fibromyalgia: a randomized trial.' *Arch Intern Med* 167(20): 2192-200.
- 142 Litz BT, Engel CC, Bryant RA, Papa A (2007). 'A randomized, controlled proof-of-concept trial of an Internet-based, therapist-assisted self management treatment for posttraumatic stress disorder.' *Am J Psychiatry* 164(11): 1676-83.
- 143 Yip YB, Sit JW, Wong DY, Chong SY, Chung LH (2008). 'A 1-year follow up of an experimental study of a self management arthritis programme with an added exercise component of clients with osteoarthritis of the knee.' *Psychol Health Med* 13(4): 402-14.
- 144 Lorig KR, Ritter PL, Laurent DD, Plant K (2008). 'The internet-based arthritis self management program: a one-year randomized trial for patients with arthritis or fibromyalgia.' *Arthritis Rheum* 59(7): 1009-17.
- 145 Deakin TA, Cade JE, Williams R, Greenwood DC (2006). 'Structured patient education: the diabetes X-PERT Programme makes a difference.' *Diabet Med* 23(9): 944-54.
- 146 Lorig KR, Ritter PL, Laurent DD, Plant K (2006). 'Internet-based chronic disease self management: a randomized trial.' *Med Care* 44(11): 964-71.
- 147 Warsi A, LaValley MP, Wang PS, et al (2003). 'Arthritis self management education programs: a meta-analysis of the effect on pain and disability.' *Arthritis Rheum* 48(8): 2207-13.
- 148 Lorig KR, Sobel DS, Ritter PL, et al (2001). 'Effect of a self management program on patients with chronic disease.' *Eff Clin Pract* 4(6): 256-62.
- 149 Clark NM, Janz NK, Dodge JA, et al (2000). 'Changes in functional health status of older women with heart disease: evaluation of a program based on self-regulation.' *J Gerontol B Psychol Sci Soc Sci* 55(2): S117-26.
- 150 Powers BJ, Olsen MK, Oddone EZ, Bosworth HB (2009). 'The effect of a hypertension self management intervention on diabetes and cholesterol control.' *Am J Med* 122(7): 639-46.
- 151 Scain SF, Friedman R, Gross JL (2009). 'A structured educational program improves metabolic control in patients with type 2 diabetes: a randomized controlled trial.' *Diabetes Educ* 35(4): 603-11.
- 152 Smeulders ES, van Haastregt JC, Ambergen T, Janssen-Boyne JJ, van Eijk JT, Kempen GI (2009). 'The impact of a self management group programme on health behaviour and healthcare utilization among congestive heart failure patients.' *Eur J Heart Fail* 11(6): 609-16.
- 153 Wattana C, Srisuphan W, Pothiban L, Upchurch SL (2007). 'Effects of a diabetes self management program on glycemic control, coronary heart disease risk, and quality of life among Thai patients with type 2 diabetes.' *Nurs Health Sci* 9(2): 135-41.
- 154 Warsi A, Wang PS, LaValley MP, et al (2004). 'Self management education programs in chronic disease: a systematic review and methodological critique of the literature.' *Arch Intern Med* 164(15): 1641-9.
- 155 Guevara JP, Wolf FM, Grum CM, Clark NM (2003). 'Effects of educational interventions for self management of asthma in children and adolescents: systematic review and meta-analysis.' *BMJ* 326(7402): 1308-9.
- 156 Parchman ML, Arambula-Solomon TG, Noel PH, Larme AC, Pugh JA (2003). 'Stage of change advancement for diabetes self management behaviors and glucose control.' *Diabetes Educ* 29(1): 128-34.
- 157 Bode C, Taal E, Emons PA, Galetzka M, Rasker JJ, Van de Laar MA (2008). 'Limited results of group self management education for rheumatoid arthritis patients and their partners: explanations from the patient perspective.' *Clin Rheumatol* 27(12): 1523-8.
- 158 Wegener ST, Mackenzie EJ, Ephraim P, Ehde D, Williams R (2009). 'Self management improves outcomes in persons with limb loss.' *Arch Phys Med Rehabil* 90(3): 373-80.
- 159 Williams GC, McGregor H, Zeldman A, Freedman ZR, Deci EL, Elder D (2005). 'Promoting glycemic control through diabetes self management: evaluating a patient activation intervention.' *Patient Educ Couns* 56(1): 28-34.
- 160 Lorig K. and Holman H (1993). 'Arthritis self management studies. A twelve year review.' *Health Ed Quart* 20: 17-28.
- 161 Warsi A, LaValley MP, Wang PS, Avorn J, Solomon DH (2003). 'Arthritis self management education programs: a meta-analysis of the effect on pain and disability.' *Arthritis Rheum* 48(8): 2207-13.
- 162 Ersek M, Turner JA, McCurry SM, Gibbons L, Kraybill BM (2003). 'Efficacy of a self management group intervention for elderly persons with chronic pain.' *Clin J Pain* 19(3): 156-67.
- 163 Seltzer MM, Litchfield LC, Kapust LR, Mayer JB (1992). 'Professional and family collaboration in case management: a hospital-based replication of a community-based study.' *Social Work in Health Care* 17(1): 1-22.
- 164 Laffel LM, Vangsness L, Connell A, et al (2003). 'Impact of ambulatory, family-focused teamwork intervention on glycemic control in youth with type 1 diabetes.' *J Pediatrics* 142(4): 409-16.
- 165 Elzen H, Slaets JP, Snijders TA, Steverink N (2007). 'Evaluation of the chronic disease self management program (CDSMP) among chronically ill older people in the Netherlands.' *Soc Sci Med* 64(9): 1832-41.
- 166 Fleissig A, Glasser B, Lloyd M (1999). 'Encouraging out-patients to make the most of their first hospital appointment: to what extent can a written prompt help patients get the information they want?' *Patient Educ Couns* 38(1): 69-79.
- 167 Perneger TV, Sudre P, Muntner P, Uldry C, Courtehouse C, Naef AF, Jacquemet S, Nicod L, Rochat T, Assal JP (2002). 'Effect of patient education on self management skills and health status in patients with asthma: a randomized trial.' *Am J Med* 113(1): 7-14.
- 168 Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM (2002). 'Self management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control.' *Diabetes Care* 25(7): 1159-71.
- 169 Figar S, Galarza C, Petrlik E, Hornstein L, Rodriguez Loria G, Waisman G, Rada M, Soriano ER, de Quiros FG (2006). 'Effect of education on blood pressure control in elderly persons: a randomized controlled trial.' *Am J Hypertens* 19(7): 737-43.
- 170 Ditewig JB, Blok H, Havers J, van Veenendaal H (2010). 'Effectiveness of self-management interventions on mortality, hospital readmissions, chronic heart failure hospitalization rate and quality of life in patients with chronic heart failure: a systematic review.' *Patient Educ Couns* 78(3): 297-315.
- 171 Gallefoss F (2004). 'The effects of patient education in COPD in a 1-year follow up randomised, controlled trial.' *Patient Educ Couns* 52(3): 259-66.
- 172 Osman LM, Calder C, Godden DJ, et al (2002). 'A randomised trial of self management planning for adult patients admitted to hospital with acute asthma.' *Thorax* 57(10): 869-74.
- 173 Groessl EJ, Cronan TA (2000). 'A cost analysis of self management programs for people with chronic illness.' *Am J Comm Psy* 28(4): 455-80.
- 174 Waxman R, Woodburn H, Powell M, Woodburn J, Blackburn S, Helliwell P (2003). 'FOOTSTEP: a randomized controlled trial investigating the clinical and cost effectiveness of a patient self management program for basic foot care in the elderly.' *J Clin Epidemiol* 56(11): 1092-9.

- 175 Schermer TR, Thoonen BP, van den Boom G, Akkermans RP, Grol RP, Folgering HT, van Weel C, van Schayck CP (2002). 'Randomized controlled economic evaluation of asthma self management in primary health care.' *Am J Respir Crit Care Med* 166(8): 1062-72.
- 176 Handley MA, Shumway M, Schillinger D (2008). 'Cost-effectiveness of automated telephone self management support with nurse care management among patients with diabetes.' *Ann Fam Med* 6(6): 512-8.
- 177 Coffman JM, Cabana MD, Halpin HA, Yelin EH (2008). 'Effects of asthma education on children's use of acute care services: a meta-analysis.' *Pediatrics* 121(3): 575-86.
- 178 Hurley MV, Walsh NE, Mitchell HL, Pimm TJ, Williamson E, Jones RH, Reeves BC, Dieppe PA, Patel A (2007). 'Economic evaluation of a rehabilitation program integrating exercise, self management, and active coping strategies for chronic knee pain.' *Arthritis Rheum* 57(7): 1220-9.
- 179 Bourbeau J, Collet JP, Schwartzman K, Ducruet T, Nault D, Bradley C (2006). 'Economic benefits of self management education in COPD.' *Chest* 130(6): 1704-11.
- 180 Gallefoss F (2010). 'The effects of patient education in COPD in a 1-year follow-up randomised, controlled trial.' *Patient Educ Couns* 52(3): 259-66.
- 181 Barlow JH, Williams B, Wright CC (1999). 'Instilling the strength to fight the pain and get on with life: learning to become an arthritis self manager through an adult education programme.' *Health Ed Res* 14(4): 533-44.
- 182 Cole JD (1998). 'Psychotherapy with the chronic pain patient using coping skills development: outcome study.' *J Occ Health Psy* 3(3): 217-26.
- 183 Lorig K, Holman HR (1989). 'Long-term outcomes of an arthritis self management study: effects of reinforcement efforts.' *Soc Sci Med* 29(2): 221-4.
- 184 Charlton I, Charlton G, Broomfield J, Mullee MA (1990). 'Evaluation of peak flow and symptoms only self management plans for control of asthma in general practice.' *BMJ* 301(6765): 1355-9.
- 185 Lorig KR, Mazonson PD, Holman HR (1993). 'Evidence suggesting that health education for self management in patients with chronic arthritis has sustained health benefits while reducing health care costs.' *Arth Rheum* 36(4): 439-46.
- 186 Fries JF, Carey C, McShane DJ (1997). 'Patient education in arthritis: randomized controlled trial of a mail-delivered program.' *J Rheumatology* 24(7): 1378-83.
- 187 Milenkovic BA, Stankovic IJ, Ilic AM, Petrovic VI (2007). 'Peak expiratory flow-guided self management treatment of asthma in Serbia.' *J Asthma* 44(9): 699-704.
- 188 Gadoury M-A, Schwartzman K, Rouleau M, Maltais F, Julien M, Beaupre A, Renzi P, Begin R, Nault D, Bourbeau J (2005). 'Self management reduces both short- and long-term hospitalisation in COPD.' *Eur Respir J* 26: 853-857.
- 189 Larson A, Ward J, Ross L, Whyatt D, Weatherston M, Landau L (2010). 'Impact of structured education and self management on rural asthma outcomes.' *Aust Fam Physician* 39(3): 141-4.
- 190 Rogers A, Robinson A, Thompson D (year unknown). *Randomised controlled trial of guided self management for patients with ulcerative colitis*. The National Primary Care Research and Development Centre. www.npcrdc.man.ac.uk/research.cfm
- 191 Robinson A, Thompson DG, Wilkin D, Roberts C, et al (2001). 'Guided self management and patient-directed follow up of ulcerative colitis: a randomised trial.' *Lancet* 358(9286): 976-81.
- 192 Kennedy A, Nelson E, Reeves D, et al (2003). 'A randomised controlled trial to assess the impact of a package comprising a patient-orientated, evidence-based self-help guidebook and patient-centred consultations on disease management and satisfaction in inflammatory bowel disease.' *Health Technol Assess* 7(28): 1-113.
- 193 Kennedy A, Reeves D, Bower P, Lee V, Middleton E, Richardson G, Gardner C, Gately C, Rogers A (2007). 'The effectiveness and cost effectiveness of a national lay-led self care support programme for patients with long-term conditions: a pragmatic randomised controlled trial.' *J Epidemiol Community Health* 61(3): 254-61.
- 194 Richardson G, Kennedy A, Reeves D, Bower P, Lee V, Middleton E, Gardner C, Gately C, Rogers A (2008). 'Cost effectiveness of the Expert Patients Programme (EPP) for patients with chronic conditions.' *J Epidemiol Community Health* 62(4): 361-7.
- 195 Wilson PM (2010). 'The UK Expert Patients Program: lessons learned and implications for cancer survivors' self-care support programs.' *J Cancer Surviv* 2(1): 45-52.
- 196 Expert Patients Programme Community Interest Company (Year unknown). *Self care reduces costs and improves health - the evidence*. London: Expert Patients Programme Community Interest Company. http://patienteducation.stanford.edu/research/Review_Findings_CDSMP_Outcomes1%208%2008.pdf
- 197 Andrade WC, Camargos P, Lasmar L, Bousquet J (2010). 'A pediatric asthma management program in a low-income setting resulting in reduced use of health service for acute asthma.' *Allergy* (published online June 2010).
- 198 Deakin T, McShane CE, Cade JE, Williams RD (2005). 'Group based training for self management strategies in people with type 2 diabetes mellitus.' *Cochrane Database Syst Rev* (2): CD003417.
- 200 Bourbeau J, Julien M, Maltais F, et al (2003). 'Reduction of hospital utilization in patients with chronic obstructive pulmonary disease: a disease-specific self management intervention.' *Arch Intern Med* 163(5): 585-91.
- 201 Lorig KR, Ritter P, Stewart A, et al (2001). 'Chronic disease self management program: 2 year health status and health care utilization outcomes.' *Medical Care* 39: 1217-23.
- 202 Gallefoss F, Bakke PS (2002). 'The effect of patient education in asthma, a randomized controlled trial.' *Tidsskr Nor Laegeforen* 122(28): 2702-6.
- 203 Krishna S, Francisco BD, Balas EA, et al (2003). 'Internet-enabled interactive multimedia asthma education program: a randomized trial.' *Pediatrics* 111(3): 503-10.
- 204 Mayo PH, Richman J, Harris HW (1990). 'Results of a program to reduce admissions for adult asthma.' *Ann Intern Med* 112(11): 864-71.
- 205 Lorig KR, Ritter PL, Gonzalez VM (2003). 'Hispanic chronic disease self management: a randomized community-based outcome trial.' *Nurs Res* 52(6): 361-9.
- 206 Wheeler JR, Janz NK, Dodge JA (2003). 'Can a disease self management program reduce health care costs? The case of older women with heart disease.' *Med Care* 41(6): 706-15.
- 207 Ghosh CS, Ravindran P, Joshi M, Stearns SC (1998). 'Reductions in hospital use from self management training for chronic asthmatics.' *Soc Sci Med* 46(8): 1087-93.
- 208 Choy DK, Tong M, Ko F, et al (1999). 'Evaluation of the efficacy of a hospital-based asthma education programme in patients of low socio-economic status in Hong Kong.' *Clin Exp Allergy* 29(1): 84-90.
- 209 Thoonen BP, Schermer TR, Van Den Boom G, Molema J, Folgering H, Akkermans RP, Grol R, Van Weel C, Van Schayck CP (2003). 'Self management of asthma in general practice, asthma control and quality of life: a randomised controlled trial.' *Thorax* 58(1): 30-6.

- 210 Gadoury MA, Schwartzman K, Rouleau M, Maltais F, Julien M, Beaupre A, Renzi P, Begin R, Nault D, Bourbeau J (2005). 'Self management reduces both short- and long-term hospitalisation in COPD.' *Eur Respir J* 26(5): 853-7.
- 211 Chiang LC, Huang JL, Yeh KW, Lu CM (2004). 'Effects of a self management asthma educational program in Taiwan based on PRECEDE-PROCEED model for parents with asthmatic children.' *J Asthma* 41(2): 205-15.
- 212 Indinnimeo L, Bonci E, Capra L, La Grutta S, Monaco F, Paravati F, Passalacqua G, Silvestre G, Duse M (2009). 'Chronic disease self management interventions might have "spill-over" effects on patients' comorbid chronic conditions.' *Pediatr Allergy Immunol* 20(7): 654-9.
- 213 Clark NM, Gong M, Kaciroti N, Yu J, Wu G, Zeng Z, Wu Z (2005). 'A trial of asthma self management in Beijing schools.' *Chronic Illn* 1(1): 31-8.
- 214 Lorig KR, Ritter P, Stewart AL, Sobel DS, Byron Jr WB, Bandura A, Gonzalez VM, Laurent DD, Holman HR (2001). 'Chronic Disease Self management Program: 2-Year Health Status and Health Care Utilization Outcomes.' *Med Care* 39(11): 1217-23.
- 215 Dunn NJ, Rehm LP, Schillaci J, Soucek J, Mehta P, Ashton CM, Yanasak E, Hamilton JD (2007). 'A randomized trial of self management and psychoeducational group therapies for comorbid chronic posttraumatic stress disorder and depressive disorder.' *J Trauma Stress* 20(3): 221-37.
- 216 Downs JA, Roberts CM, Blackmore AM, Le Souëf PN, Jenkins SC (2006). 'Benefits of an education programme on the self management of aerosol and airway clearance treatments for children with cystic fibrosis.' *Chron Respir Dis* 3(1): 19-27.
- 217 Wetzels R, van Weel C, Grol R, Wensing M (2008). 'Family practice nurses supporting self management in older patients with mild osteoarthritis: a randomized trial.' *BMC Fam Pract* 9: 7.
- 218 Buszewicz M, Rait G, Griffin M, Nazareth I, Patel A, Atkinson A, Barlow J, Haines A (2006). 'Self management of arthritis in primary care: randomised controlled trial.' *BMJ* 333(7574): 879.
- 219 Jowett S, Bryan S, Murray E, McCahon D, Raftery J, Hobbs FD, Fitzmaurice D (2006). 'Patient self management of anticoagulation therapy: a trial-based cost-effectiveness analysis.' *Br J Haematol* 134(6): 632-9.
- 220 Monninkhof E, van der Valk P, Schermer T, van der Palen J, van Herwaarden C, Zielhuis G (2004). 'Economic evaluation of a comprehensive self management programme in patients with moderate to severe chronic obstructive pulmonary disease.' *Chron Respir Dis* 1(1): 7-16.
- 221 Buszewicz M, Rait G, Griffin M, Nazareth I, Patel A, Atkinson A, Barlow J, Haines A (2006). 'Self management of arthritis in primary care: randomised controlled trial.' *BMJ* 333: 879.
- 222 Monninkhof E, van der Valk P, van der Palen J, et al (2003). 'Self management education for patients with chronic obstructive pulmonary disease: a systematic review.' *Thorax* 58(5): 394-8.
- 223 Barlow, Wright, Bancroft, Turner (2003). *Evaluation of the Chronic Disease Self management Course*. UK Department of Health and Long-term Medical Conditions Alliance.
- 224 Patel A, Buszewicz M, Beecham J, Griffin M, Rait G, Nazareth I, Atkinson A, Barlow J, Haines A (2009). 'Economic evaluation of arthritis self management in primary care.' *BMJ* 339: b3532.
- 225 McPherson AC, Glazebrook C, Forster D, James C, Smyth A (2006). 'A randomized, controlled trial of an interactive educational computer package for children with asthma.' *Pediatrics* 117(4): 1046-54.
- 226 Foster G, Taylor SJ, Eldridge SE, Ramsay J, Griffiths CJ (2007). 'Self management education programmes by lay leaders for people with chronic conditions.' *Cochrane Database Syst Rev* (4): CD005108.
- 227 Glasgow NJ, Ponsonby AL, Yates R, et al (2003). 'Proactive asthma care in childhood: general practice based randomised controlled trial.' *BMJ* 327(7416):659.
- 228 Feldman PH, Peng TR, Murtaugh CM, Kelleher C, Donelson SM, McCann ME, Putnam ME (2004). 'A randomized intervention to improve heart failure outcomes in community-based home health care.' *Home Health Care Serv Q* 23(1): 1-23.
- 229 Morrison A (2001). 'Effectiveness of printed patient educational materials in chronic illness: a systematic review of controlled trials.' *J Manag Pharm Care* 1(1): 51-62.
- 230 Gibson PG, Powell H, Coughlan J et al (2004). 'Limited (information only) patient education programs for adults with asthma (Cochrane Review).' *The Cochrane Library*, Issue 2. Chichester: John Wiley & Sons.
- 231 Kennedy A, Robinson A, Hann M et al (2003). 'A cluster-randomised controlled trial of a patient-centred guidebook for patients with ulcerative colitis: effect on knowledge, anxiety and quality of life.' *Health Soc Care Community* 11(1): 64-72.
- 232 Lancaster T, Stead LF. Self-help interventions for smoking cessation. *Cochrane Database of Systematic Reviews* 2005, Issue 3.
- 233 Protheroe J, Blakeman T, Bower P, Chew-Graham C, Kennedy A (2010). 'An intervention to promote patient participation and self-management in long term conditions: development and feasibility testing.' *BMC Health Serv Res* 10: 206.
- 234 Maly RC, Abrahamse AF, Hirsch SH et al (1996). 'What influences physician practice behavior? An interview study of physicians who received consultative geriatric assessment recommendations.' *Arch Family Med*; 5: 448-454.
- 235 Laffel LM, Vangsness L, Connell A et al (2003). 'Impact of ambulatory, family-focused teamwork intervention on glycemic control in youth with type 1 diabetes.' *J Pediatrics* 142(4): 409-16.
- 236 Estabrooks C, Goel V, Thiel E et al (2001). 'Decision aids: are they worth it? A systematic review.' *J Health Serv Res Policy* 6(3): 170-82.
- 237 Sturt J, Taylor H, Docherty A et al (2006). 'A psychological approach to providing self management education for people with type 2 diabetes: the Diabetes Manual.' *BMC Fam Pract* 7: 70.
- 238 Robinson T (2010). 'Developing self management plans to help people with COPD to control their condition.' *Nurs Times* 106(17): 14.
- 239 Flesher M, Woo P, Chiu A, Charlebois A, Warburton DE, Leslie B (2010). 'Self-management and biomedical outcomes of a cooking, and exercise program for patients with chronic kidney disease.' *J Ren Nutr* (published online July 2010).
- 240 Piette JD, Rosland AM, Silveira M, Kabeto M, Langa KM (2010). 'The case for involving adult children outside of the household in the self-management support of older adults with chronic illnesses.' *Chronic Illn* 6(1): 34-45.
- 241 Rosland AM, Heisler M, Choi HJ, Silveira MJ, Piette JD (2010). 'Family influences on self-management among functionally independent adults with diabetes or heart failure: do family members hinder as much as they help?' *Chronic Illn* 6(1): 22-33.
- 242 Hibbard JH, Mahoney E (2010). 'Toward a theory of patient and consumer activation.' *Patient Educ Couns* 78(3):377-81.
- 243 Remmers C, Hibbard J, Mosen DM, Wagenfield M, Hoyer RE, Jones C (2009). 'Is patient activation associated with future health outcomes and healthcare utilization among patients with diabetes?' *J Ambul Care Manage* 32(4):320-7.

- 244 Cunningham AJ, Lockwood GA, Cunningham JA (1991). 'A relationship between perceived self-efficacy and quality of life in cancer patients.' *Patient Educ Couns* 17(1):71-8.
- 245 Hibbard JH, Greene J, Tusler M (2009). 'Improving the outcomes of disease management by tailoring care to the patient's level of activation.' *Am J Manag Care* 15(6):353-60.
- 246 Wagner EH, LaCroix AZ, Grothaus L et al (1994). 'Preventing disability and falls in older adults: A population-based randomized trial.' *Am J Public Health* 84(11): 1800-6.
- 247 Roberts J, Browne GB, Streiner D (1995). 'Problem-solving counselling or phone-call support for outpatients with chronic illness: effective for whom?' *Can J Nurs Res* 27(3): 111-37.
- 248 Gani F, Pozzi E, Crivellaro MA, et al (2001). 'The role of patient training in the management of seasonal rhinitis and asthma: clinical implications.' *Allergy* 56(1): 65-8.
- 249 Benton JM, O'Hara PA, Chen H et al (1997). 'Changing bowel hygiene practice successfully.' *Geriatr Nurs* 18(1): 12-7.
- 250 Krumholz HM, Amatruda J, Smith GL (2002). 'Randomized trial of an education and support intervention to prevent readmission of patients with heart failure.' *J Am Coll Cardio* 39(1): 83-9.
- 251 Ruetsch C (2010). 'Practice strategies to improve compliance and patient self-management.' *J Manag Care Pharm* 16(1 Suppl B): S26-7.
- 252 The British Liver Trust (1999). *Living a Healthy Life with Long-term Illness*. Leland Stanford Junior University.
- 253 Barlow JH, Turner, Wright (1998). 'Sharing, caring and learning to take control: self management training for people with arthritis.' *Psy, Health Med* 3(4): 387-93.
- 254 Barlow JH, Turner, Wright (1998). 'Long-term outcomes of an arthritis self management programme.' *Brit J Rheumatology* 37: 1315-9.
- 255 Barlow JH, Turner, Wright (2000). 'A randomised controlled study of the arthritis self management programme in the UK.' *Health Ed Res* 15(6): 665-80.
- 256 Barlow JH, Turner, Wright (2001). 'Patient education for people with arthritis in rural communities: the UK experience.' *Patient Ed Couns* 44: 205-14.
- 257 Barlow JH, Wright CC, Sheasby et al (2002). 'Self management approaches for people with chronic conditions: a review.' *Patient Ed Couns* 48: 177-87.
- 258 Department of Health (2001). *The Expert Patient: A New Approach to Chronic Disease Management for the 21st Century*. London: Department of Health.
- 259 Brown SA. Studies of educational interventions and outcomes in diabetic adults: a meta-analysis revisited. *Patient Educ Couns* 1990; 16(3): 189-215.
- 260 Guevara JP, Wolf FM, Grum CM, Clark NM (2003). 'Effects of educational interventions for self management of asthma in children and adolescents: systematic review and meta-analysis.' *BMJ* 326(7402): 1308-9.
- 261 Norris SL, Lau J, Smith SJ, et al (2002). 'Self management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control.' *Diabetes Care* 25(7): 1159-71.
- 262 Lorig K. and Holman H (1993). 'Arthritis self management studies. A twelve year review.' *Health Ed Quart* 20: 17-28.
- 263 Warsi A, LaValley MP, Wang PS, et al (2003). 'Arthritis self management education programs: a meta-analysis of the effect on pain and disability.' *Arthritis Rheum* 48(8): 2207-13.
- 264 Reid MC, Papaleontiou M, Ong A, Breckman R, Wethington E, Pillemer K (2008). 'Self management strategies to reduce pain and improve function among older adults in community settings: a review of the evidence.' *Pain Med* 9(4): 409-24.
- 265 Bodenheimer T, Lorig K, Holman H, Grumbach K (2002). 'Patient self management of chronic disease in primary care.' *JAMA* 288(19): 2469-75.
- 266 Yu PL, Ye W, Liu XR, et al (2003). 'Evaluation on the effectiveness for self management of hypertensive patients in a community.' *Zhonghua Liu Xing Bing Xue Za Zhi* 24(9): 790-3.
- 267 Wright SP, Walsh H, Ingley KM et al (2003). 'Uptake of self management strategies in a heart failure management programme.' *Eur J Heart Fail* 5(3): 371-80.
- 268 Lorig K, Holman HR (1989). 'Long-term outcomes of an arthritis self management study: effects of reinforcement efforts.' *Soc Sci Med* 29(2): 221-4.
- 269 Charlton I, Charlton G, Broomfield J, Mullee MA (1990). 'Evaluation of peak flow and symptoms only self management plans for control of asthma in general practice.' *BMJ* 301(6765): 1355-9.
- 270 Lorig KR, Mazonson PD, Holman HR (1993). 'Evidence suggesting that health education for self management in patients with chronic arthritis has sustained health benefits while reducing health care costs.' *Arth Rheum* 36(4): 439-46.
- 271 Fries JF, Carey C, McShane DJ (1997). 'Patient education in arthritis: randomized controlled trial of a mail-delivered program.' *J Rheumatology* 24(7): 1378-83.
- 272 Rogers A, Robinson A, Thompson D (Year unknown). *Randomised controlled trial of guided self management for patients with ulcerative colitis*. The National Primary Care Research and Development Centre. www.npcrdc.man.ac.uk/research.cfm
- 273 Robinson A, Thompson DG, Wilkin D, Roberts C et al (2001). 'Guided self management and patient-directed follow up of ulcerative colitis: a randomised trial.' *Lancet* 358(9286): 976-81.
- 274 Gallefoss F (2004). 'The effects of patient education in COPD in a 1-year follow up randomised, controlled trial.' *Patient Educ Couns* 52(3): 259-66.
- 275 Lorig KR, Ritter P, Stewart A et al (2001). 'Chronic disease self management program: 2 year health status and health care utilization outcomes.' *Medical Care* 39: 1217-23.
- 276 Gallefoss F, Bakke PS (2002). 'Cost-benefit and cost-effectiveness analysis of self management in patients with COPD - a 1 year follow up randomized, controlled trial.' *Respir Med* 96(6): 424-31.
- 277 Bradley C, Blenkinsopp A. Over the counter drugs. The future for self medication. *BMJ* 1996; 312 (7034): 835-837.
- 278 Garcia-Alamino JM, Ward AM, Alonso-Coello P, Perera R, Bankhead C, Fitzmaurice D, Heneghan CJ (2010). 'Self-monitoring and self-management of oral anticoagulation.' *Cochrane Database Syst Rev* 4: CD003839.
- 279 Powers BJ, Trinh JV, Bosworth HB (2010). 'Can this patient read and understand written health information?' *JAMA* 304(1): 76-84.
- 280 Wensing M, Vingerhoets E, Grol R (2003). 'Feedback based on patient evaluations: a tool for quality improvement?' *Patient Educ Couns* 51(2): 149-53.
- 281 Glasgow RE, Strycker LA, King DK, Toobert DJ, Rahm AK, Jex M, Nutting PA (2006). 'Robustness of a computer-assisted diabetes self management intervention across patient characteristics, healthcare settings, and intervention staff.' *Am J Manag Care* 12(3): 137-45.
- 282 Ruetsch C (2010). 'Practice strategies to improve compliance and patient self-management.' *J Manag Care Pharm* 16(1 Suppl B): S26-7.
- 283 Carryer J, Budge C, Hansen C, Gibbs K (2010). 'Providing and receiving self-management support for chronic illness: patients' and health practitioners' assessments.' *J Prim Health Care* 2(2): 124-9.

- 284 Siminerio LM, Funnell MM, Peyrot M, Rubin RR (2010). 'US nurses' perceptions of their role in diabetes care: results of the cross-national Diabetes Attitudes Wishes and Needs (DAWN) study.' *Diabetes Educ* 33(1): 152-62.
- 285 Williams GC, McGregor HA, King D, Nelson CC, Glasgow RE (2010). 'Variation in perceived competence, glycemic control, and patient satisfaction: relationship to autonomy support from physicians.' *Patient Educ Couns* 57(1): 39-45.
- 286 Lam TP, Lam KF (2010). 'Family doctors' attitudes towards patient self-management of upper respiratory tract infections.' *Hong Kong Med J* 7(2): 146-9.
- 287 Steurer-Stey C, Fletcher M, Vetter W, Steurer J (2010). 'Patient education in asthma: a survey of physicians' knowledge of the principles and implementation of self management in practice.' *Swiss Med Wkly* 136(35-36): 561-5.
- 288 McIntosh A, Shaw CF (2010). 'Barriers to patient information provision in primary care: patients' and general practitioners' experiences and expectations of information for low back pain.' *Health Expect* 6(1): 19-29.
- 289 Koura MR, Khairy AE, Abdel-Aal NM, Mohamed HF, Amin GA, Sabra AY (2010). 'The role of primary health care in patient education for diabetes control.' *J Egypt Public Health Assoc* 76(3-4): 241-64.
- 290 Puder JJ, Keller U (2010). 'Quality of diabetes care: problem of patient or doctor adherence?' *Swiss Med Wkly* 133(39-40): 530-4.
- 291 Washburn SC, Hornberger CA, Klutman A, Skinner L (2010). 'Nurses' knowledge of heart failure education topics as reported in a small midwestern community hospital.' *J Cardiovasc Nurs* 20(3): 215-20.
- 292 Wetzels R, Geest TA, Wensing M et al (2004). 'GPs' views on involvement of older patients: an European qualitative study.' *Patient Educ Couns* 53(2): 183-8.
- 293 Rogers A, Kennedy A, Nelson E, Robinson A (2005). 'Uncovering the limits of patient-centeredness: implementing a self management trial for chronic illness.' *Qual Health Res* 15(2): 224-39.
- 294 Kennedy A, Gask L, Rogers A (2005). 'Training professionals to engage with and promote self management.' *Health Educ Res* 20(5): 567-78.
- 295 Wolters R, Wensing M, Van Weel C, Grol R (2004). 'The effect of a distance-learning programme on patient self management of Lower Urinary Tract Symptoms (LUTS) in general practice: a randomised controlled trial.' *Eur Urol* 46(1): 95-101.
- 296 Ding D, Liu HY, Cooper R, Cooper RA, Smailagic A, Siewiorek D (2010). 'Virtual coach technology for supporting self-care.' *Phys Med Rehabil Clin N Am* 21(1): 179-94.
- 297 Britt E, Blampied NM (2010). 'Motivational interviewing training: a pilot study of the effects on practitioner and patient behaviour.' *Behav Cogn Psychother* 38(2): 239-44.
- 298 de Ridder DT, Theunissen NC, van Dulmen SM (2010). 'Does training general practitioners to elicit patients' illness representations and action plans influence their communication as a whole?' *Patient Educ Couns* 66(3): 327-36.
- 299 Blackstien-Hirsch P, Anderson G, Cicutto L, Mclvor A, Norton P (2010). 'Implementing continuing education strategies for family physicians to enhance asthma patients' quality of life.' *J Asthma* 37(3): 247-57.
- 300 Lacroix A, Assal JP (2010). 'Is it possible to improve the teaching methods of physicians who instruct patients? Comparative analysis of the same courses given before and after teacher's training.' *Diabete Metab* 18(5): 387-94.
- 301 Kosmala-Anderson JP, Wallace LM, Turner A (2010). 'Confidence matters: a Self-Determination Theory study of factors determining engagement in self-management support practices of UK clinicians.' *Psychol Health Med* 15(4): 478-91.
- 302 Cohen D, Longo MF, Hood K, et al (2004). 'Resource effects of training general practitioners in risk communication skills and shared decision making competences.' *J Eval Clin Pract* 10(3): 439-45.
- 303 Edwards A, Elwyn G, Hood K, et al (2004). 'Patient-based outcome results from a cluster randomized trial of shared decision making skill development and use of risk communication aids in general practice.' *Fam Pract* 21(4): 347-54.
- 304 Cockle-Hearne J, Faithfull S (2010). 'Self-management for men surviving prostate cancer: a review of behavioural and psychosocial interventions to understand what strategies can work, for whom and in what circumstances.' *Psychooncology* 19(9): 909-22.
- 305 Smeulders ES, van Haastregt JC, Ambergen T, Uszko-Lencer NH, Janssen-Boyne JJ, Gorgels AP, Stoffers HE, Lodewijks-van der Bolt CL, van Eijk JT, Kempen GI (2010). 'Nurse-led self-management group programme for patients with congestive heart failure: randomized controlled trial.' *J Adv Nurs* 66(7): 1487-99.
- 306 D'Eramo Melkus G, Chyun D, Vorderstrasse A, Newlin K, Jefferson V, Langerman S (2010). 'The effect of a diabetes education, coping skills training, and care intervention on physiological and psychosocial outcomes in black women with type 2 diabetes.' *Biol Res Nurs* 12(1): 7-19.
- 307 Armitage CJ, Arden MA (2010). 'How useful are the stages of change for targeting interventions? Randomized test of a brief intervention to reduce smoking.' *Health Psychol* 27(6): 789-98.
- 308 Tzeng LF, Chiang LC, Hsueh KC, Ma WF, Fu LS (2010). 'A preliminary study to evaluate a patient-centred asthma education programme on parental control of home environment and asthma signs and symptoms in children with moderate-to-severe asthma.' *J Clin Nurs* 19(9-10): 1424-33.
- 309 Druss BG, Zhao L, von Esenwein SA, Bona JR, Fricks L, Jenkins-Tucker S, Sterling E, Diclemente R, Lorig K (2010). 'The Health and Recovery Peer (HARP) Program: a peer-led intervention to improve medical self-management for persons with serious mental illness.' *Schizophr Res* 118(1-3): 264-70.
- 310 Funnell MM (2010). 'Peer-based behavioural strategies to improve chronic disease self-management and clinical outcomes: evidence, logistics, evaluation considerations and needs for future research.' *Fam Pract* 27(Suppl 1): i 17-22.
- 311 New N (2010). 'Teaching so they hear: using a co-created diabetes self-management education approach.' *J Am Acad Nurse Pract* 22(6): 316-25.
- 312 Dorn SD (2010). 'Systematic review: self-management support interventions for irritable bowel syndrome.' *Aliment Pharmacol Ther* 32(4): 513-21.
- 313 Iversen MD, Hammond A, Betteridge N (2010). 'Self-management of rheumatic diseases: state of the art and future perspectives.' *Ann Rheum Dis* 69(6): 955-63.
- 314 Fors T, Damsgård E, Roe C, Anke A (2010). 'Readiness to adopt a self-management approach to pain - Are profiles of subscale scores on the Pain Stages of Change Questionnaire useful?' *Eur J Pain* (published online May 2010).
- 315 Coffman JM, Cabana MD, Halpin HA, Yelin EH (2008). 'Effects of asthma education on children's use of acute care services: a meta-analysis.' *Pediatrics* 121(3): 575-86.
- 316 Shigaki CL, Smarr KL, Yang Gong, Donovan-Hanson K, Siva C, Johnson RA, Ge B, Musser DR (2008). 'Social interactions in an online self management program for rheumatoid arthritis.' *Chronic Illn* 4(4): 239-46.
- 317 Goudswaard AN, Stolk RP, Zuihoff NP, de Valk HW, Rutten GE (2004). 'Long-term effects of self management education for patients with Type 2 diabetes taking maximal oral hypoglycaemic therapy: a randomized trial in primary care.' *Diabet Med* 21(5): 491-6.

- 318 Schneider S, Iannotti RJ, Nansel TR, Haynie DL, Simons-Morton B, Sobel DO, Zeitoff L, Clark L, Plotnick LP (2007). 'Identification of distinct self management styles of adolescents with type 1 diabetes.' *Diabetes Care* 30(5): 1107-12.
- 319 Osborne RH, Buchbinder R, Ackerman IN (2006). 'Can a disease-specific education program augment self management skills and improve Health-Related Quality of Life in people with hip or knee osteoarthritis?' *BMC Musculoskelet Disord* 7: 90.
- 320 Damush TM, Wu J, Bair MJ, Sutherland JM, Kroenke K (2008). 'Self management practices among primary care patients with musculoskeletal pain and depression.' *J Behav Med* 31(4): 301-7.
- 321 Jerant A, Kravitz R, Moore-Hill M, Franks P (2008). 'Depressive symptoms moderated the effect of chronic illness self management training on self-efficacy.' *Med Care* 46(5): 523-31.
- 322 Hoffmann T, McKenna K, Worrall L, Read SJ (2007). 'Randomised trial of a computer-generated tailored written education package for patients following stroke.' *Age Ageing* 36(3): 280-6.
- 323 Thoolen B, de Ridder D, Bensing J, Gorter K, Rutten G (2008). 'Beyond Good Intentions: the development and evaluation of a proactive self management course for patients recently diagnosed with type 2 diabetes.' *Health Educ Res* 23(1): 53-61.
- 324 King DK, Estabrooks PA, Strycker LA, Toobert DJ, Bull SS, Glasgow RE (2006). 'Outcomes of a multifaceted physical activity regimen as part of a diabetes self management intervention.' *Ann Behav Med* 31(2): 128-37.
- 325 DeWalt DA, Malone RM, Bryant ME, Kosnar MC, Corr KE, Rothman RL, Sueta CA, Pignone MP (2006). 'A heart failure self management program for patients of all literacy levels: a randomized, controlled trial.' *BMC Health Serv Res* 6: 30.
- 326 Nguyen HQ, Carrieri-Kohlman V (2005). 'Dyspnea self management in patients with chronic obstructive pulmonary disease: moderating effects of depressed mood.' *Psychosomatics* 46(5): 402-10.
- 327 North Simcoe Muskoka Chronic Disease Prevention and Management Regional Action Group (2009). *Directions from a Local Scan: Self management and Empowering the Person Living with Diabetes in the North Simcoe Muskoka LHIN*. North Simcoe Muskoka Chronic Disease Prevention and Management Regional Action Group.
- 328 *Preventing and Managing Chronic Disease: Ontario's Framework*.
- 329 Weng LC, Dai YT, Huang HL, Chiang YJ (2010). 'Self-efficacy, self-care behaviours and quality of life of kidney transplant recipients.' *J Adv Nurs* 66(4): 828-38.
- 330 Brewer-Lowry AN, Arcury TA, Bell RA, Quandt SA (2010). 'Differentiating Approaches to Diabetes Self-Management of Multi-ethnic Rural Older Adults at the Extremes of Glycemic Control.' *Gerontologist* 50(5): 657-67.
- 331 Auslander WF, Sterzing PR, Zayas LE, White NH (2010). 'Psychosocial resources and barriers to self-management in African American adolescents with type 2 diabetes: a qualitative analysis.' *Diabetes Educ* 36(4): 613-22.
- 332 Crowe M, Whitehead L, Jo Gagan M, Baxter D, Panckhurst A (2010). 'Self-management and chronic low back pain: a qualitative study.' *J Adv Nurs* 66(7): 1478-86.
- 333 Merelle SY, Sorbi MJ, Duivenvoorden HJ, Passchier J (2010). 'Qualities and health of lay trainers with migraine for behavioral attack prevention.' *Headache* 50(4): 613-25.
- 334 Given CW, Given BA, Sikorskii A, You M, Jeon S, Champion V, McCorkle R (2010). 'Deconstruction of nurse-delivered patient self-management interventions for symptom management: factors related to delivery enactment and response.' *Ann Behav Med* 40(1): 99-113.
- 335 Buccheri RK, Trygstad LN, Buffum MD, Lyttle K, Dowling G (2010). 'Comprehensive evidence-based program teaching self-management of auditory hallucinations on inpatient psychiatric units.' *Issues Ment Health Nurs* 31(3): 223-31.
- 336 Pitt VJ, O'connor D, Green S (2010). 'Referral of people with osteoarthritis to self-management programmes: barriers and enablers identified by general practitioners.' *Disabil Rehabil* 30(25): 1938-46.
- 337 Kosmala-Anderson JP, Wallace LM, Turner A (2010). 'Confidence matters: a Self-Determination Theory study of factors determining engagement in self-management support practices of UK clinicians.' *Psychol Health Med* 15(4): 478-91.
- 338 Audulv A, Asplund K, Norbergh KG (2010). 'Who's in charge? The role of responsibility attribution in self-management among people with chronic illness.' *Patient Educ Couns* 81(1): 94-100.
- 339 Hibbard JH, Collins PA, Mahoney E, Baker LH (2010). 'The development and testing of a measure assessing clinician beliefs about patient self-management.' *Health Expect* 13(1): 65-72.
- 340 Blakeman T, Bower P, Reeves D, Chew-Graham C (2010). 'Bringing self-management into clinical view: a qualitative study of long-term condition management in primary care consultations.' *Chronic Illn* 6(2): 136-50.
- 341 Bodenheimer T, MacGregor K, Shafiri C (2005). *Helping Patients Manage Their Chronic Conditions*. California: California Healthcare Foundation.
- 342 Shirazi M, Parikh SV, Alaeddini F, Lonka K, Zeinaloo AA, Sadeghi M, Arbabi M, Nejatiasafa AA, Shahrivar Z, Wahlstrom R (2010). 'Effects on knowledge and attitudes of using stages of change to train general practitioners on management of depression: a randomized controlled study.' *Can J Psychiatry* 54(10): 693-700.
- 343 Dennis SM, Zwar N, Griffiths R, et al. Chronic disease management in primary care: from evidence to policy. *Med J Aust* 2008; 188 (8 Suppl): S53-S56.
- 344 Jordan JE, Osborne RH. Chronic disease self management education programs: challenges ahead. *Med J Aust* 2007; 186: 84-87.
- 345 Newman S, Steed L, Mulligan K. Self management interventions for chronic illness. *Lancet* 2004; 364: 1523-1537.
- 346 Anderson L. et al (2005). 'Self-help books for depression: how can practitioners and patients make the right choice?' *Br J Gen Pract* 55 (514): 387-392.
- 347 Glasgow NJ, Ponsonby AL, Yates R, et al (2003). 'Proactive asthma care in childhood: general practice based randomised controlled trial.' *BMJ* 327(7416): 659.
- 348 Little P, Dorward M, Warner G et al (2004). 'Randomised controlled trial of effect of leaflets to empower patients in consultations in primary care.' *BMJ* 328(7437): 441.
- 349 Fleissig A, Glasser B, Lloyd M (1999). 'Encouraging out-patients to make the most of their first hospital appointment: to what extent can a written prompt help patients get the information they want?' *Patient Educ Couns* 38(1): 69-79.
- 350 Dally DL, Dahar W, Scott A, et al (2002). 'The impact of a health education program targeting patients with high visit rates in a managed care organization.' *Am J Health Promot* 17(2): 101-11.
- 351 van Boeijen CA et al (2005). 'Efficacy of self-help manuals for anxiety disorders in primary care: a systematic review.' *Fam Pract* 22(2): 192-196.
- 352 Roberts L, Little P, Chapman J, Cantrell T, Pickering R, Langridge J (2010). 'The back home trial: general practitioner-supported leaflets may change back pain behavior.' *Spine* 27(17): 1821-8.
- 353 Morrison A (2001). 'Effectiveness of printed patient educational materials in chronic illness: a systematic review of controlled trials.' *J Manag Pharm Care* 1(1): 51-62.

- 354 Gibson PG, Powell H, Coughlan J et al (2004). 'Limited (information only) patient education programs for adults with asthma (Cochrane Review).' *The Cochrane Library*, Issue 2. Chichester: John Wiley & Sons.
- 355 Lorig KR, Ritter PL, Laurent DD, Fries JF (2004). 'Long-term randomized controlled trials of tailored-print and small-group arthritis self management interventions.' *Med Care* 42(4): 346-54.
- 356 Forster A, Smith J, Young J et al (2004). 'Information provision for stroke patients and their caregivers (Cochrane Review).' *The Cochrane Library*, Issue 2. Chichester: John Wiley & Sons.
- 357 Seals TD, Keith MR (1997). 'Influence of patient information leaflets on anticonvulsant drug compliance in prison.' *Am J Health Syst Pharm* 54(22): 2585-7.
- 358 Kennedy A, Robinson A, Hann M et al (2003). 'A cluster-randomised controlled trial of a patient-centred guidebook for patients with ulcerative colitis: effect on knowledge, anxiety and quality of life.' *Health Soc Care Community* 11(1): 64-72.
- 359 Lafata JE, Baker AM, Divine GW et al (2002). 'The use of computerized birthday greeting reminders in the management of diabetes.' *J Gen Intern Med* 17(7): 521-30.
- 360 Sethares KA, Elliott K (2004). 'The effect of a tailored message intervention on heart failure readmission rates, quality of life, and benefit and barrier beliefs in persons with heart failure.' *Heart Lung* 33(4): 249-60.
- 361 Enwald HP, Huotari ML (2010). 'Preventing the obesity epidemic by second generation tailored health communication: an interdisciplinary review.' *J Med Internet Res* 12(2): e24.
- 362 Osman LM, Abdalla MI, Beattie JAG et al (1994). 'Reducing hospital admission through computer supported education for asthma patients.' *BMJ* 308(6928): 568-71.
- 363 Azrin NH, Teichner G (1998). 'Evaluation of an instructional program for improving medication compliance for chronically mentally ill outpatients.' *Behav Res Ther* 36(9): 849-61.
- 364 Oermann MH, Webb SA, Ashare JA (2003). 'Outcomes of videotape instruction in clinic waiting area.' *Orthopaedic Nurs* 22(2): 102-5.
- 365 Grilli R, Ramsay C, Minozzi S (2004). 'Mass media interventions: effects on health services utilisation (Cochrane Review).' *The Cochrane Library* (Issue 2). Chichester: John Wiley & Sons.
- 366 Buchbinder R (2008). 'Self management education en masse: effectiveness of the Back Pain: Don't Take It Lying Down mass media campaign.' *Med J Aust* 189(10 Suppl): S29-S32.
- 367 Williams A, Hagerty BM, Brasington SJ, Clem JB, Williams DA (2010). 'Stress Gym: Feasibility of deploying a web-enhanced behavioral self-management program for stress in a military setting.' *Mil Med* 175(7): 487-93.
- 368 Cordina M, McElnay JC, Hughes CM (2001). 'Assessment of a community pharmacy-based program for patients with asthma.' *Pharmacotherapy* 21(10): 1196-203.
- 369 Samoocha D, Bruinvels DJ, Elbers NA, Anema JR, van der Beek AJ (2010). 'Effectiveness of web-based interventions on patient empowerment: a systematic review and meta-analysis.' *J Med Internet Res* 12(2): e23.
- 370 Rybarczyk B, DeMarco G, DeLaCruz M, Lapidus S (1999). 'Comparing mind-body wellness interventions for older adults with chronic illness: classroom versus home instruction.' *Beh Med* 24(4): 181-90.
- 371 Schaffer SD, Tian L (2004). 'Promoting adherence: effects of theory-based asthma education.' *Clin Nurs Res* 13(1): 69-89.
- 372 Shames RS, Sharek P, Mayer M, et al (2004). 'Effectiveness of a multicomponent self management program in at-risk, school-aged children with asthma.' *Ann Allergy Asthma Immunol* 92(6): 611-8.
- 373 Krishna S, Francisco BD, Balas EA et al (2003). 'Internet-enabled interactive multimedia asthma education program: a randomized trial.' *Pediatrics* 111(3): 503-10.
- 374 Glasgow RE, Nutting PA, Toobert DJ et al (2006). 'Effects of a brief computer-assisted diabetes self management intervention on dietary, biological and quality-of-life outcomes.' *Chronic Illn* 2(1): 27-38.
- 375 Yeh YT, Chiu YT, Liu CT et al (2006). 'Development and evaluation of an integrated patient-oriented education management system for diabetes.' *Stud Health Technol Inform* 122: 172-5.
- 376 Savage I, Goodyer L (2003). 'Providing information on metered dose inhaler technique: is multimedia as effective as print?' *Fam Pract* 20(5): 552-7.
- 377 Bowen AM, Horvath K, Williams ML (2007). 'A randomized control trial of Internet-delivered HIV prevention targeting rural MSM.' *Health Educ Res* 22(1): 120-7.
- 378 De Bourdeaudhuij I, Stevens V, Vandelanotte C, Brug J (2007). 'Evaluation of an interactive computer-tailored nutrition intervention in a real-life setting.' *Ann Behav Med* 33(1): 39-48.
- 379 Proudfoot J, Parker G, Hyett M, Manicavasagar V, Smith M, Grdovic S, Greenfield L (2007). 'Next generation of self management education: Web-based bipolar disorder program.' *Aust N Z J Psychiatry* 41(11): 903-9.
- 380 Gerber BS, Brodsky IG, Lawless KA, Smolin LI, Arozullah AM, Smith EV, Berbaum ML, Heckerling PS, Eiser AR (2005). 'Implementation and evaluation of a low-literacy diabetes education computer multimedia application.' *Diabetes Care* 28(7): 1574-80.
- 381 van der Meer V, Bakker MJ, van den Hout WB, Rabe KF, Sterk PJ, Kievit J, Assendelft WJ, Sont JK (2009). 'Internet-based self management plus education compared with usual care in asthma: a randomized trial.' *Ann Intern Med* 151(2): 110-20.
- 382 Nguyen HQ, Donesky-Cuenco D, Wolpin S, Reinke LF, Benditt JO, Paul SM, Carrieri-Kohlman V (2008). 'Randomized controlled trial of an internet-based versus face-to-face dyspnea self management program for patients with chronic obstructive pulmonary disease: pilot study.' *J Med Internet Res* 10(2): e9.
- 383 Bingham CR, Barretto AI, Walton MA, Bryant CM, Shope JT, Raghunathan TE (2010). 'Efficacy of a web-based, tailored, alcohol prevention/intervention program for college students: initial findings.' *J Am Coll Health* 58(4): 349-56.
- 384 Krishna S, Balas EA, Spencer DC, et al (1997). 'Clinical trials of interactive computerized patient education: implications for family practice.' *J Fam Pract* 45(1): 25-33.
- 385 Stinson J, Wilson R, Gill N, Yamada J, Holt J (2009). 'A systematic review of internet-based self management interventions for youth with health conditions.' *J Pediatr Psychol* 34(5): 495-510.
- 386 Mulvaney SA, Rothman RL, Wallston KA, Lybarger C, Dietrich MS (2010). 'An internet-based program to improve self-management in adolescents with type 1 diabetes.' *Diabetes Care* 33(3): 602-4.
- 387 Jernigan VB, Lorig K (2010). 'The Internet Diabetes Self-Management Workshop for American Indians and Alaska Natives.' *Health Promot Pract* (published online June 2010).
- 388 Ghahari S, Leigh Packer T, Passmore AE (2010). 'Effectiveness of an online fatigue self-management programme for people with chronic neurological conditions: a randomized controlled trial.' *Clin Rehabil* 24(8): 727-44.

- 389 Lorig K, Ritter PL, Laurent DD, Plant K, Green M, Jernigan VB, Case S (2010). 'Online diabetes self-management program: a randomized study.' *Diabetes Care* 33(6): 1275-81.
- 390 Wanner M, Martin-Diener E, Braun-Fahrländer C, Bauer G, Martin BW (2010). 'Effectiveness of active-online, an individually tailored physical activity intervention, in a real-life setting: randomized controlled trial.' *J Med Internet Res* 11(3): e23.
- 391 Eysenbach G, Powell J, Englesakis M et al (2004). 'Health related virtual communities and electronic support groups: systematic review of the effects of online peer to peer interactions.' *BMJ* 328(7449): 1166.
- 392 Hoffman-Goetz L, Donelle L (2007). 'Chat room computer-mediated support on health issues for aboriginal women.' *Health Care Women Int* 28(4): 397-418.
- 393 Barrera M Jr, Glasgow RE, McKay HG, Boles SM, Feil EG (2002). 'Do Internet-based support interventions change perceptions of social support?: An experimental trial of approaches for supporting diabetes self management.' *Am J Community Psychol* 30(5): 637-54.
- 394 Stinson JN, McGrath PJ, Hodnett ED, Feldman BM, Duffy CM, Huber AM, Tucker LB, Hetherington CR, Tse SM, Spiegel LR, Campillo S, Gill NK, White ME (2010). 'An internet-based self-management program with telephone support for adolescents with arthritis: a pilot randomized controlled trial.' *J Rheumatol* 37(9): 1944-52.
- 395 Simoni JM, Huh D, Frick PA, Pearson CR, Andrasik MP, Dunbar PJ, Hooton TM (2009). 'Peer support and pager messaging to promote antiretroviral modifying therapy in Seattle: a randomized controlled trial.' *J Acquir Immune Defic Syndr* 52(4): 465-473.
- 396 Faridi Z, Liberti L, Shuval K, Northrup V, Ali A, Katz DL (2008). 'Evaluating the impact of mobile telephone technology on type 2 diabetic patients' self management: the NICHE pilot study.' *J Eval Clin Pract* 14(3): 465-9.
- 397 Franklin VL, Waller A, Pagliari C, Greene SA (2006). 'A randomized controlled trial of Sweet Talk, a text-messaging system to support young people with diabetes.' *Diabet Med* 23(12): 1332-8.
- 398 Plotnikoff RC, McCargar LJ, Wilson PM, Loucaides CA (2005). 'Efficacy of an e-mail intervention for the promotion of physical activity and nutrition behavior in the workplace context.' *Am J Health Promot* 19(6): 422-9.
- 399 Yap TL, Hemmings A, Davis LS (2010). 'The systematic development of a tailored e-mail intervention for health behavior change toward increasing intentional physical activity.' *West J Nurs Res* 31(3): 330-46.
- 400 Ersser SJ, Cowdell FC, Latter SM, Healy E (2010). 'Self-management experiences in adults with mild-moderate psoriasis: an exploratory study and implications for improved support.' *Br J Dermatol* (published online June 2010).
- 401 Seymour CJ, Cannon S (2010). 'Student-facilitated health promotion intervention for chronic disease self-management in at-risk elders reflections from the field.' *J Allied Health* 39(2): 120-8.
- 402 Morice AH, Wrench C (2001). 'The role of the asthma nurse in treatment compliance and self management following hospital admission.' *Respir Med* 95(11): 851-6.
- 403 Duke SA, Colagiuri S, Colagiuri R (2009). 'Individual patient education for people with type 2 diabetes mellitus.' *Cochrane Database Syst Rev* (1): CD005268.
- 404 Bailey WC, Richards JM Jr, Brooks CM (1990). 'A randomized trial to improve self management practices of adults with asthma.' *Arch Intern Med* 150(8): 1664-8.
- 405 Boulware LE, Daumit GL, Frick KD et al (2001). 'An evidence-based review of patient-centered behavioral interventions for hypertension.' *Am J Prev Med* 21(3): 221-32.
- 406 Dracup K, Baker DW, Dunbar SB, et al (1994). 'Management of heart failure. II. Counseling, education, and lifestyle modifications.' *JAMA* 272(18): 1442-6.
- 407 Superio-Cabuslay E, Ward MM, Lorig KR (1996). 'Patient education interventions in osteoarthritis and rheumatoid arthritis: a meta-analytic comparison with nonsteroidal antiinflammatory drug treatment.' *Arthritis Care Res* 9(4): 292-301.
- 408 Goudswaard AN, Stolk RP, Zuithoff NP, et al (2004). 'Long-term effects of self management education for patients with Type 2 diabetes taking maximal oral hypoglycaemic therapy: a randomized trial in primary care.' *Diabet Med* 21(5): 491-6.
- 409 Evans R 3rd, Gergen PJ, Mitchell H et al (1999). 'A randomized clinical trial to reduce asthma morbidity among inner-city children: results of the National Cooperative Inner-City Asthma Study.' *J Pediatr* 135(3): 332-8.
- 410 Lawn S, Battersby MW, Pols RG et al (2007). 'The mental health expert patient: findings from a pilot study of a generic chronic condition self management programme for people with mental illness.' *Int J Soc Psychiatry* 53(1): 63-74.
- 411 Lee JK, Grace KA, Taylor AJ (2006). 'Effect of a pharmacy care program on medication adherence and persistence, blood pressure, and low-density lipoprotein cholesterol: a randomized controlled trial.' *JAMA* 296(21): 2563-71.
- 412 Gaston MH, Porter GK, Thomas VG (2007). 'Prime Time Sister Circles: evaluating a gender-specific, culturally relevant health intervention to decrease major risk factors in mid-life African-American women.' *J Natl Med Assoc* 99(4): 428-38.
- 413 Bordeleau L et al (2003). 'Quality of life in a randomized trial of group psychosocial support in metastatic breast cancer: overall effects of the intervention and an exploration of missing data.' *J Clin Oncol* 21(10): 1944-1951.
- 414 Lorig KR, Ritter PL, Gonzalez VM (2003). 'Hispanic chronic disease self management: a randomized community-based outcome trial.' *Nurs Res* 52(6): 361-9.
- 415 Carone M, Bertolotti G, Cerveri I, De Benedetto F, Fogliani V, Nardini S, Portalone L, Rossi A, Sanguinetti CM, Schiavina M, Donner CF (2002). 'EDU-CARE, a randomised, multicentre, parallel group study on education and quality of life in COPD.' *Monaldi Arch Chest Dis* 57(1): 25-9.
- 416 Swerissen H, Belfrage J, Weeks A, Jordan L, Walker C, Furler J, McAvoy B, Carter M, Peterson C (2006). 'A randomised control trial of a self management program for people with a chronic illness from Vietnamese, Chinese, Italian and Greek backgrounds.' *Patient Educ Couns* 64(1-3): 360-8.
- 417 Lorig K, Ritter PL, Plant K (2005). 'A disease-specific self-help program compared with a generalized chronic disease self-help program for arthritis patients.' *Arthritis Rheum* 53(6): 950-7.
- 418 Griffiths C, Motlib J, Azad A, Ramsay J, Eldridge S, Feder G, Khanam R, Munni R, Garrett M, Turner A, Barlow J (2005). 'Randomised controlled trial of a lay-led self management programme for Bangladeshi patients with chronic disease.' *Br J Gen Pract* 55(520): 831-7.
- 419 Schiel R, Braun A, Muller R, Helbich C, Siefke S, Franke I, Osterbrink B, Leppert K, Stein G, Muller UA (2004). 'A structured treatment and educational program for patients with type 2 diabetes mellitus, insulin therapy and impaired cognitive function (DikoL).' *Med Klin* 99(6): 285-92.
- 420 Battersby M, Hoffmann S, Cadilhac D, Osborne R, Lalor E, Lindley R (2009). 'Getting your life back on track after stroke': a Phase II multi-centered, single-blind, randomized, controlled trial of the Stroke Self management Program vs. the Stanford Chronic Condition Self management Program or standard care in stroke survivors.' *Int J Stroke* 4(2): 137-44.

- 421 The EPP Evaluation Team (2005). *Process Evaluation of the EPP – Report II: Examination of the implementation of the Expert Patients Programme within the structures and locality contexts of the NHS in England*. University of Manchester, National Primary Care Research and Development Centre.
- 422 Wallace AS, Carlson JR, Malone RM, Joyner J, Dewalt DA (2010). 'The influence of literacy on patient-reported experiences of diabetes self-management support.' *Nurs Res* 59(5): 356-63.
- 423 Greenhalgh T, Campbell-Richards D, Vijayaraghavan S, Collard A, Malik F, Griffin M, Morris J, Claydon A, Macfarlane F (2010). 'New models of self-management education for minority ethnic groups: pilot randomized trial of a story-sharing intervention.' *J Health Serv Res Policy* (published online August 2010).
- 424 Walker EA, Stevens KA, Persaud S (2010). 'Promoting diabetes self-management among African Americans: an educational intervention.' *J Health Care Poor Underserved* 21(3 Suppl): 169-86.
- 425 Press VG, Pincavage AT, Pappalardo AA, Baker DC, Conwell WD, Cohen JC, Hoyte FL, Johnson ME, Prochaska MH, Vela MB, Arora VM (2010). 'The Chicago Breathe Project: a regional approach to improving education on asthma inhalers for resident physicians and minority patients.' *J Natl Med Assoc* 102(7): 548-55.
- 426 Coleman S, McQuade J, Rose J, Inderjeeth C, Carroll G, Briffa NK (2010). 'Self-management for osteoarthritis of the knee: does mode of delivery influence outcome?' *BMC Musculoskelet Disord* 11: 56.
- 427 Schillinger D, Hammer H, Wang F, Palacios J, McLean I, Tang A, Youmans S, Handley M (2010). 'Seeing in 3-D: examining the reach of diabetes self-management support strategies in a public health care system.' *Health Educ Behav* 35(5): 664-82.
- 428 Department of Health (2007). *Self care Support: The evidence pack – summary of work in progress 2005-07*. London: Department of Health.
- 429 Campbell HS et al (2004). 'Cancer peer support programs - do they work?' *Patient Ed Counsel* 55: 3-15.
- 430 Corben S, Rosen R (2005). *Self management for long-term conditions*. London: King's Fund.
- 431 van Hooff ML, van der Merwe JD, O'Dowd J, Pavlov PW, Spruit M, de Kleuver M, van Limbeek J (2010). 'Daily functioning and self-management in patients with chronic low back pain after an intensive cognitive behavioral programme for pain management.' *Eur Spine J* 19(9): 1517-26.
- 432 Clark NM, Gong M, Kaciroti N et al (2005). 'A trial of asthma self management in Beijing schools.' *Chronic Illn* 1(1): 31-8.
- 433 Heisler M (2006). *Building Peer Support Programmes to Manage Disease: Seven Models for Success*. California Health Care Foundation.
- 434 Chien WT et al (2004). 'A randomized controlled trial of a mutual support group for family caregivers of patients with schizophrenia.' *Int J Nurs Stud* 41(6): 637-649.
- 435 Lorig KR, Sobel DS, Stewart AL, et al (1999). 'Evidence suggesting that a chronic disease self management program can improve health status while reducing utilization and costs: A randomized trial.' *Med Care* 37(1): 5-14.
- 436 Warsi A, Wang PS, LaValley MP, et al (2004). 'Self management education programs in chronic disease: a systematic review and methodological critique of the literature.' *Arch Intern Med* 164(15): 1641-9.
- 437 Bernard-Bonnin AC, Stachenko S, Bonin D, et al (1995). 'Self management teaching programs and morbidity of pediatric asthma: a meta-analysis.' *J Allergy Clin Immunol* 95(1 Pt 1): 34-41.
- 438 Monninkhof E, van der Valk P, van der Palen J et al (2003). 'Self management education for patients with chronic obstructive pulmonary disease: a systematic review.' *Thorax* 58(5): 394-8.
- 439 Kennedy A, Reeves D, Bower P et al (2007). 'The effectiveness and cost effectiveness of a national lay-led self care support programme for patients with long-term conditions: a pragmatic randomised controlled trial.' *J Epidemiol Community Health* 61(3): 254-61.
- 440 Monninkhof EM, et al (2003). 'Self management education for chronic obstructive pulmonary disease.' *Cochrane Database Syst Rev* (1): CD002990.
- 441 Boulware LE et al (2001). 'An evidence-based review of patient-centered behavioural interventions for hypertension.' *Am J Prev Med* 21(3): 221-232.
- 442 Niedermann K et al (2004). 'Gap between short- and long-term effects of patient education in rheumatoid arthritis patients: a systematic review.' *Arthritis Rheum* 51(3): 388-398.
- 443 Ersek M, Turner JA, Cain KC, Kemp CA (2008). 'Results of a randomized controlled trial to examine the efficacy of a chronic pain self management group for older adults.' *Pain* 138(1): 29-40.
- 444 Chodosh J et al (2005). 'Meta-analysis: chronic disease self management programs for older adults.' *Ann Intern Med* 143(6): 427-438.
- 445 Cooper H et al (2001). 'Chronic disease patient education: lessons from meta-analyses.' *Patient Educ Couns* 44 (2): 107-117.
- 446 Banet GA, Felchli MA (1997). 'The potential utility of a shared medical record in a "first-time" stroke population.' *J Vasc Nurs* 15(1): 29-33.
- 447 Warner JP, King M, Blizard R, et al (2000). 'Patient-held shared care records for individuals with mental illness. Randomised controlled evaluation.' *Br J Psychiatry* 177: 319-24.
- 448 Lester H, Allan T, Wilson S, Jowett S, Roberts L (2003). 'A cluster randomised controlled trial of patient-held medical records for people with schizophrenia receiving shared care.' *Br J Gen Pract* 53(488): 197-203.
- 449 Ball MJ, Smith C, Bakalar RS (2007). 'Personal health records: empowering consumers.' *J Healthc Inf Manag* 21(1): 76-86.
- 450 Winkelman WJ, Leonard KJ, Rossos PG (2005). 'Patient-perceived usefulness of online electronic medical records: employing grounded theory in the development of information and communication technologies for use by patients living with chronic illness.' *J Am Med Inform Assoc* 12(3): 306-14.
- 451 Currell R, Urquhart C (2004). 'Nursing record systems: effects on nursing practice and health care outcomes (Cochrane Review).' *The Cochrane Library* (Issue 2).
- 452 Ross SE, Moore LA, Earnest MA et al (2004). 'Providing a web-based online medical record with electronic communication capabilities to patients with congestive heart failure: randomized trial.' *J Med Internet Res* 6(2): e12.
- 453 Estabrooks C, Goel V, Thiel E et al (2001). 'Decision aids: are they worth it? A systematic review.' *J Health Serv Res Policy* 6(3): 170-82.
- 454 Schillinger D, Hammer H, Wang F, Palacios J, McLean I, Tang A, Youmans S, Handley M (2008). 'Seeing in 3-D: examining the reach of diabetes self management support strategies in a public health care system.' *Health Educ Behav* 35(5): 664-82.
- 455 Smith SR, Rublein JC, Marcus C, Brock TP, Chesney MA (2003). 'A medication self management program to improve adherence to HIV therapy regimens.' *Patient Educ Couns* 50(2): 187-99.
- 456 McGeoch GR, Willsman KJ, Dowson CA, Town GI, Frampton CM, McCartin FJ, Cook JM, Epton MJ (2006). 'Self management plans in the primary care of patients with chronic obstructive pulmonary disease.' *Respirology* 11(5): 611-8.

- 457 Garrett N, Hageman CM, Sibley SD, Davern M, Berger M, Brunzell C, Malecha K, Richards SW (2005). 'The effectiveness of an interactive small group diabetes intervention in improving knowledge, feeling of control, and behavior.' *Health Promot Pract* 6(3): 320-8.
- 458 Kaya Z, Erkan F, Ozkan M, Ozkan S, Kocaman N, Ertekin BA, Direk N (2009). 'Self management plans for asthma control and predictors of patient compliance.' *J Asthma* 46(3): 270-5.
- 459 Janson SL, McGrath KW, Covington JK, Cheng SC, Boushey HA (2009). 'Individualized asthma self management improves medication adherence and markers of asthma control.' *J Allergy Clin Immunol* 123(4): 840-6.
- 460 Magar Y, Vervloet D, Steenhouwer F, Smaga S, Mechin H, Rocca Serra JP, Marchand C, d'Ivernois JF (2005). 'Assessment of a therapeutic education programme for asthma patients: "un souffle nouveau."' *Patient Educ Couns* 58(1): 41-6.
- 461 D'Souza WJ, Te Karu H, Fox C, Harper M, Gemmell T, Ngatuere M, Wickens K, Crane J, Pearce N, Beasley R (1998). 'Long-term reduction in asthma morbidity following an asthma self management programme.' *ERS J*.
- 462 Gallefoss F, Bakke PS (1999). 'How does patient education and self management among asthmatics and patients with chronic obstructive pulmonary disease affect medication?' *Am J Respir Crit Care Med* 160(6): 2000-5.
- 463 Gillies J, Barry D, Crane J, et al (1996). 'A community trial of a written self management plan for children with asthma.' *NZ Med J* 109(1015): 30-3.
- 464 van der Palen J, Klein JJ, Zielhuis GA, van Herwaarden CL (1998). 'The role of self-treatment guidelines in self management education for adult asthmatics.' *Respir Med* 92(4): 668-75.
- 465 Starnino VR, Mariscal S, Holter MC, Davidson LJ, Cook KS, Fukui S, Rapp CA (2010). 'Outcomes of an illness self-management group using wellness recovery action planning.' *Psychiatr Rehabil J* 34(1): 57-60.
- 466 Burkhart PV, Rayens MK, Oakley MG, Abshire DA, Zhang M (2007). 'Testing an intervention to promote children's adherence to asthma self management.' *J Nurs Scholarsh* 39(2): 133-40.
- 467 Kennedy AP, Nelson E, Reeves D, Richardson G, Roberts C, Robinson A, Rogers AE, Sculpher M, Thompson DG (2004). 'A randomised controlled trial to assess the effectiveness and cost of a patient orientated self management approach to chronic inflammatory bowel disease.' *Gut* 53(11): 1639-45.
- 468 Milenkovic B, Bosnjak-Petrovic V (2007). 'Self management program in treatment of asthma.' *Srp Arh Celok Lek* 135(3-4): 147-52.
- 469 Powell H, Gibson PG (2004). 'Options for self management education for adults with asthma (Cochrane Review).' *The Cochrane Library* (Issue 2). Chichester: John Wiley & Sons.
- 470 Gibson PG, Powell H, Coughlan J, Wilson AJ, Abramson M, Haywood P, Bauman A, Hensley MJ, Walters EH (2003). 'Self management education and regular practitioner review for adults with asthma.' *Cochrane Database Syst Rev* (1): CD001117.
- 471 Gallefoss F, Bakke PS (1999). 'How does patient education and self management among asthmatics and patients with chronic obstructive pulmonary disease affect medication?' *Am J Respir Crit Care Med* 160(6): 2000-5.
- 472 Gillies J, Barry D, Crane J, et al (1996). 'A community trial of a written self management plan for children with asthma.' *NZ Med J* 109(1015): 30-3.
- 473 Jones A, Pill R, Adams S (2010). 'Qualitative study of views of health professionals and patients on guided self management plans for asthma.' *BMJ* 321(7275): 1507-10.
- 474 Daly JM, Buckwalter K, Maas M. Written and computerized care plans (2002). 'Organizational processes and effect on patient outcomes.' *J Gerontol Nurs* 28(9): 14-23.
- 475 Toelle BG, Ram FSF (2004). 'Written individualised management plans for asthma in children and adults (Cochrane Review).' *The Cochrane Library* (Issue 2). Chichester: John Wiley & Sons.
- 476 van der Palen J, Klein JJ, Zielhuis GA, van Herwaarden CL (1998). 'The role of self-treatment guidelines in self management education for adult asthmatics.' *Respir Med* 92(4): 668-75.
- 477 Couturaud F, Proust A, Frachon I, Dewitte JD, Oger E, Quiot JJ, Leroyer C (2002). 'Education and self management: a one-year randomized trial in stable adult asthmatic patients.' *J Asthma* 39(6): 493-500.
- 478 Walters JA, Turnock AC, Walters EH, Wood-Baker R (2010). 'Action plans with limited patient education only for exacerbations of chronic obstructive pulmonary disease.' *Cochrane Database Syst Rev* 5: CD005074.
- 479 Gibson PG, Powell H, Coughlan J, et al (2004). 'Self management education and regular practitioner review for adults with asthma (Cochrane Review).' *The Cochrane Library* (Issue 2). Chichester: John Wiley & Sons.
- 480 Gallefoss F (2004). 'The effects of patient education in COPD in a 1-year follow up randomised, controlled trial.' *Patient Educ Couns* 52(3): 259-66.
- 481 Espinoza-Palma T, Zamorano A, Arancibia F, Bustos MF, Silva MJ, Cardenas C, De La Barra P, Puente V, Cerda J, Castro-Rodriguez JA, Prado F (2009). 'Effectiveness of asthma education with and without a self management plan in hospitalized children.' *J Asthma* 46(9): 906-10.
- 482 Wood-Baker R, McGlone S, Venn A, Walters EH (2006). 'Written action plans in chronic obstructive pulmonary disease increase appropriate treatment for acute exacerbations.' *Respirology* 11(5): 619-26.
- 483 van Eijken M, Wensing M, de Konink M, Vernooij M, Zielhuis G, Lagro T, Rikkert MO, Grol R (2004). 'Health education on self management and seeking health care in older adults: a randomised trial.' *Patient Educ Couns* 55(1): 48-54.
- 484 Shepperd S, Parkes J, McClaren J, Phillips C (2004). 'Discharge planning from hospital to home (Cochrane Review).' *The Cochrane Library* (Issue 2). Chichester: John Wiley & Sons.
- 485 Spillane LL, Lumb EW, Cobaugh DJ, et al (1997). 'Frequent users of the emergency department: can we intervene?' *Acad Emerg Med* 4(6): 574-80.
- 486 Brown MD, Reeves MJ, Meyerson K, Korzeniewski SJ (2006). 'Randomized trial of a comprehensive asthma education program after an emergency department visit.' *Ann Allergy Asthma Immunol* 97(1): 44-51.
- 487 Osman LM, Calder C, Godden DJ, Friend JA, McKenzie L, Legge JS, Douglas JG (2002). 'A randomised trial of self management planning for adult patients admitted to hospital with acute asthma.' *Thorax* 57(10): 869-74.
- 488 Ersek M, Turner JA, Cain KC, Kemp CA (2004). 'Chronic pain self management for older adults: a randomized controlled trial.' *BMC Geriatr* 4: 7.
- 489 Barlow JH, Turner AP, Gilchrist M (2009). 'A randomised controlled trial of lay-led self management for myocardial infarction patients who have completed cardiac rehabilitation.' *Eur J Cardiovasc Nurs* 8(4): 293-301.
- 490 Kline KS, Scott LD, Britton AS (2007). 'The use of supportive-educative and mutual goal-setting strategies to improve self management for patients with heart failure.' *Home Healthc Nurse* 25(8): 502-10.
- 491 Brown CT, Yap T, Cromwell DA, Rixon L, Steed L, Mulligan K, Mundy A, Newman SP, van der Meulen J, Emberton M (2007). 'Self management for men with lower urinary tract symptoms: randomised controlled trial.' *BMJ* 334(7583): 25.

- 492 Glasgow RE, Toobert DJ, Hampson SE, Strycker LA (2002). 'Implementation, generalization and long-term results of the "choosing well" diabetes self management intervention.' *Patient Educ Couns* 48(2): 115-22.
- 493 Goeppinger J, Lorig KR, Ritter PL, Mutatkar S, Villa F, Gizlice Z (2009). 'Mail-delivered arthritis self management tool kit: a randomized trial and longitudinal followup.' *Arthritis Rheum* 61(7): 867-75.
- 494 Braun AK, Kubiak T, Kuntsche J, Meier-Hofig M, Muller UA, Feucht I, Zeyfang A (2009). 'SGS: a structured treatment and teaching programme for older patients with diabetes mellitus - a prospective randomised controlled multi-centre trial.' *Age Ageing* 38(4): 390-6.
- 495 Bastiaenen CH, de Bie RA, Vlaeyen JW, Goossens ME, Leffers P, Wolters PM, Bastiaanssen JM, Brandt PA, Essed GG (2008). 'Long-term effectiveness and costs of a brief self management intervention in women with pregnancy-related low back pain after delivery.' *BMC Pregnancy Childbirth* 8: 19.
- 496 Scholz U, Knoll N, Sniehotta FF, Schwarzer R (2006). 'Physical activity and depressive symptoms in cardiac rehabilitation: long-term effects of a self management intervention.' *Soc Sci Med* 62(12): 3109-20.
- 497 Estabrooks PA, Nelson CC, Xu S, King D, Bayliss EA, Gaglio B, Nutting PA, Glasgow RE (2005). 'The frequency and behavioral outcomes of goal choices in the self management of diabetes.' *Diabetes Educ* 31(3): 391-400.
- 498 Brody BL, Roch-Levecq AC, Thomas RG, Kaplan RM, Brown SI (2005). 'Self management of age-related macular degeneration at the 6-month follow up: a randomized controlled trial.' *Arch Ophthalmol* 123(1): 46-53.
- 499 Thoolen B, de Ridder D, Bensing J, Gorter K, Rutten G (2008). 'Beyond Good Intentions: the development and evaluation of a proactive self management course for patients recently diagnosed with type 2 diabetes.' *Health Educ Res* 23(1): 53-61.
- 500 Garrett DG, Bluml BM (2005). 'Patient self management program for diabetes: first-year clinical, humanistic, and economic outcomes.' *J Am Pharm Assoc* 45: 130-137.
- 501 West SP, Laguna C, Trief PM, Izquierdo R, Weinstock RS (2010). 'Goal setting using telemedicine in rural underserved older adults with diabetes: experiences from the informatics for diabetes education and telemedicine project.' *Telemed J E Health* 16(4): 405-16.
- 502 Russell CL (2010). 'A clinical nurse specialist-led intervention to enhance medication adherence using the plan-do-check-act cycle for continuous self-improvement.' *Clin Nurse Spec* 24(2): 69-75.
- 503 Wheeler JR, Janz NK, Dodge JA (2003). 'Can a disease self management program reduce health care costs? The case of older women with heart disease.' *Med Care* 41(6): 706-15.
- 504 Janson SL, Fahy JV, Covington JK, Paul SM, Gold WM, Boushey HA (2003). 'Effects of individual self management education on clinical, biological, and adherence outcomes in asthma.' *Am J Med* 115(8): 620-6.
- 505 Lerman I, Lopez-Ponce A, Villa AR, Escobedo M, Caballero EA, Velasco ML, Gomez-Perez FJ, Rull-Rodrigo JA (2009). 'Pilot study of two different strategies to reinforce self care behaviors and treatment compliance among type 2 diabetes patients from low income strata.' *Gac Med Mex* 145(1): 15-9.
- 506 Anderson RM, Funnell MM, Nwankwo R, Gillard ML, Oh M, Fitzgerald JT (2005). 'Evaluating a problem-based empowerment program for African Americans with diabetes: results of a randomized controlled trial.' *Ethn Dis* 15(4): 671-8.
- 507 Shearer NB, Cisar N, Greenberg EA (2007). 'A telephone-delivered empowerment intervention with patients diagnosed with heart failure.' *Heart Lung* 36(3): 159-69.
- 508 Glasgow RE, Nutting PA, Toobert DJ, King DK, Strycker LA, Jex M, O'Neill C, Whitesides H, Merenich J (2006). 'Effects of a brief computer-assisted diabetes self management intervention on dietary, biological and quality-of-life outcomes.' *Chronic Illn* 2(1): 27-38.
- 509 Blixen CE, Bramstedt KA, Hammel JP, Tilley BC. A pilot study of health education via a nurse-run telephone self management programme for elderly people with osteoarthritis. *J Telemed Telecare* 2004; 10(1): 44-9.
- 510 Heaney CA, Goetzel RZ (1997). 'A review of health-related outcomes of multi-component worksite health promotion programs.' *Am J Health Promot* 11(4): 290-307.
- 511 Riley KM, Glasgow RE, Eakin EG (2001). 'Resources for health: A social-ecological intervention for supporting self management of chronic conditions.' *J Health Psy* 6(6): 693-705.
- 512 Clark M, Hampson SE, Avery L, Simpson R (2004). 'Effects of a tailored lifestyle self management intervention in patients with type 2 diabetes.' *Br J Health Psychol* 9(Pt 3): 365-79.
- 513 Samuel-Hodge CD, Keyserling TC, Park S, Johnston LF, Gizlice Z, Bangdiwala SI (2009). 'A randomized trial of a church-based diabetes self management program for African Americans with type 2 diabetes.' *Diabetes Educ* 35(3): 439-54.
- 514 Bourbeau J, Julien M, Maltais F, Rouleau M, Beupre A, Begin R, Renzi P, Nault D, Borycki E, Schwartzman K, Singh R, Collet JP (2003). 'Reduction of hospital utilization in patients with chronic obstructive pulmonary disease: a disease-specific self management intervention.' *Arch Intern Med* 163(5): 585-91.
- 515 Glasgow RE, La Chance PA, Toobert DJ, et al (1997). 'Long-term effects and costs of brief behavioural dietary intervention for patients with diabetes delivered from the medical office.' *Patient Educ Couns* 32(3): 175-84.
- 516 Blanson Henkemans OA, van der Boog PJ, Lindenberg J, van der Mast CA, Neerincx MA, Zwetsloot-Schonk BJ (2009). 'An online lifestyle diary with a persuasive computer assistant providing feedback on self management.' *Technol Health Care* 17(3): 253-67.
- 517 Monninkhof E, van der Valk P, van der Palen J, van Herwaarden C, Zielhuis G (2003). 'Effects of a comprehensive self management programme in patients with chronic obstructive pulmonary disease.' *Eur Respir J* 22(5): 815-20.
- 518 Brown SA, Blozis SA, Kouzekanani K, Garcia AA, Winchell M, Hanis CL (2005). 'Dosage effects of diabetes self management education for Mexican Americans: the Starr County Border Health Initiative.' *Diabetes Care* 28(3): 527-32.
- 519 Hendricks LE, Hendricks RT (2000). 'The effect of diabetes self management education with frequent follow up on the health outcomes of African American men.' *Diabetes Educ* 26(6): 995-1002.
- 520 Young D, Furler J, Vale M, et al (2007). 'Patient Engagement and Coaching for Health: The PEACH study - a cluster randomised controlled trial using the telephone to coach people with type 2 diabetes to engage with their GPs to improve diabetes care.' *BMC Fam Pract* 8: 20.
- 521 LaFramboise LM, Toder CM, Zimmerman L, Agrawal S (2003). 'Comparison of Health Buddy with traditional approaches to heart failure management.' *Fam Community Health* 26(4): 275-88.
- 522 Dilorio C, Reisinger EL, Yeager KA, McCarty F (2009). 'A telephone-based self management program for people with epilepsy.' *Epilepsy Behav* 14(1): 232-6.
- 523 Esters ON, Boeckner LS, Hubert M, Horacek T, Kritsch KR, Oakland MJ, Lohse B, Greene G, Nitzke S (2010). 'Educator and participant perceptions and cost analysis of stage-tailored educational telephone calls.' *J Nutr Educ Behav* 40(4): 258-64.

- 524 Kim HS, Oh JA (2003). 'Adherence to diabetes control recommendations: impact of nurse telephone calls.' *J Adv Nurs* 44(3): 256-61.
- 525 Whitlock WL, Brown A, Moore K, et al (2000). 'Telemedicine improved diabetic management.' *Mil Med* 165(8): 579-84.
- 526 Shearer NB, Cisar N, Greenberg EA (2007). 'A telephone-delivered empowerment intervention with patients diagnosed with heart failure.' *Heart Lung* 36(3): 159-69.
- 527 Piette JD, Weinberger M, Kraemer FB, McPhee SJ (2001). 'Impact of automated calls with nurse follow up on diabetes treatment outcomes in a Department of Veterans Affairs Health Care System: a randomized controlled trial.' *Diabetes Care* 24(2): 202-8.
- 528 West SP, Laguna C, Trief PM, Izquierdo R, Weinstock RS (2010). 'Goal setting using telemedicine in rural underserved older adults with diabetes: experiences from the informatics for diabetes education and telemedicine project.' *Telemed J E Health* 16(4): 405-16.
- 529 Piette JD, Weinberger M, McPhee SJ, et al (2000). 'Do automated calls with nurse follow up improve self care and glycemic control among vulnerable patients with diabetes?' *Am J Med* 108(1): 20-7.
- 530 Oh JA, Kim HS, Yoon KH, Choi ES (2003). 'A telephone-delivered intervention to improve glycemic control in type 2 diabetic patients.' *Yonsei Med J* 44(1):1-8.
- 531 Linden A, Butterworth SW, Prochaska JO (2010). 'Motivational interviewing-based health coaching as a chronic care intervention.' *J Eval Clin Pract* 16(1): 166-74.
- 532 Paradis V, Cossette S, Frasure-Smith N, Heppell S, Guertin MC (2010). 'The efficacy of a motivational nursing intervention based on the stages of change on self-care in heart failure patients.' *J Cardiovasc Nurs* 25(2): 130-41.
- 533 Oslin DW, Sayers S, Ross J, et al (2003). 'Disease management for depression and at-risk drinking via telephone in an older population of veterans.' *Psychosom Med* 65(6): 931-7.
- 534 Riegel B, Carlson B, Kopp Z, et al (2002). 'Effect of a standardized nurse case-management telephone intervention on resource use in patients with chronic heart failure.' *Arch Intern Med* 162(6): 705-12.
- 535 Jarrett ME, Cain KC, Burr RL, Hertig VL, Rosen SN, Heitkemper MM (2009). 'Comprehensive self management for irritable bowel syndrome: randomized trial of in-person vs. combined in-person and telephone sessions.' *Am J Gastroenterol* 104(12): 3004-14.
- 536 Given CW, Bradley C, You M, Sikorskii A, Given B (2010). 'Costs of novel symptom management interventions and their impact on hospitalizations.' *J Pain Symptom Manage* 39(4): 663-72.
- 537 Anderson DR, Christison-Lagay J, Villagra V, Liu H, Dziura J (2010). 'Managing the Space between Visits: A Randomized Trial of Disease Management for Diabetes in a Community Health Center.' *J Gen Intern Med* (published online June 2010).
- 538 Hopp FP, Hogan MM, Woodbridge PA, Lowery JC (2010). 'The use of telehealth for diabetes management: a qualitative study of telehealth provider perceptions.' *Implement Sci* 2: 14.
- 539 Faghri PD, Blozie E, Gustavesen S, Kotejoshyer R (2010). 'The role of tailored consultation following health-risk appraisals in employees' health behavior.' *J Occup Environ Med* 50(12): 1378-85.
- 540 Sorensen G, Barbeau EM, Stoddard AM, et al (2007). 'Tools for health: the efficacy of a tailored intervention targeted for construction laborers.' *Cancer Causes Control* 18(1): 51-9.
- 541 Aldana SG, Greenlaw RL, Diehl HA, et al (2005). 'The effects of a worksite chronic disease prevention program.' *J Occup Environ Med* 47(6): 558-64.
- 542 Proper KI, van der Beek AJ, Hildebrandt VH, et al (2004). 'Worksite health promotion using individual counselling and the effectiveness on sick leave; results of a randomised controlled trial.' *Occup Environ Med* 61(3): 275-9.
- 543 Muto T, Yamauchi K (2001). 'Evaluation of a multicomponent workplace health promotion program conducted in Japan for improving employees' cardiovascular disease risk factors.' *Prev Med* 33(6): 571-7.
- 544 Bloch MJ, Armstrong DS, Dettling L, et al (2006). 'Partners in lowering cholesterol: comparison of a multidisciplinary educational program, monetary incentives, or usual care in the treatment of dyslipidemia identified among employees.' *J Occup Environ Med* 48(7): 675-81.
- 545 Golaszewski T, Yen LT (1992). 'Demographic characteristics of users of worksite health promotion written materials.' *Am J Health Promot* 6(5): 359-63.
- 546 Anderson DR, Stauffer MJ (1996). 'The impact of worksite-based health risk appraisal on health-related outcomes: a review of the literature.' *Am J Health Promot* 10(6): 499-508.
- 547 Tessaro IA, Taylor S, Belton L, et al (2000). 'Adapting a natural (lay) helpers model of change for worksite health promotion for women.' *Health Educ Res* 15(5): 603-14.
- 548 Fujii H, Nakade M, Haruyama Y, Fukuda H, Hashimoto M, Ikuyama T, Kaburagi H, Murai E, Okumura M, Sairenchi T, Muto T (2010). 'Evaluation of a computer-tailored lifestyle modification support tool for employees in Japan.' *Ind Health* 47(3): 333-41.
- 549 Gates D, Brehm B, Hutton S, et al (2006). 'Changing the work environment to promote wellness: a focus group study.' *AAOHN J* 54(12): 515-20.
- 550 Heaney CA, Goetzel RZ (1997). 'A review of health-related outcomes of multi-component worksite health promotion programs.' *Am J Health Promot* 11(4): 290-307.
- 551 Pelletier KR (2001). 'A review and analysis of the clinical and cost-effectiveness studies of comprehensive health promotion and disease management programs at the worksite: 1998-2000 update.' *Am J Health Promot* 16(2): 107-16.
- 552 Bradley C, Blenkinsopp A (1996). 'Over the counter drugs. The future for self medication.' *BMJ* 312 (7034): 835-837.
- 553 Coster S, et al (2000). 'Monitoring blood glucose control in diabetes mellitus: a systematic review.' *Health Technology Assessment* 4 (12).
- 554 Jansen JP (2006). 'Self-monitoring of glucose in type 2 diabetes mellitus: a Bayesian meta-analysis of direct and indirect comparisons.' *Curr Med Res Opin* 22(4): 671-681.
- 555 Fitzmaurice DA, Murray ET, Gee KM, Allan TF, Hobbs FD (2002). 'A randomised controlled trial of patient self management of oral anticoagulation treatment compared with primary care management.' *J Clin Pathol* 55(11): 845-9.
- 556 Bennett H, Laird K, Margolius D, Ngo V, Thom DH, Bodenheimer T (2009). 'The effectiveness of health coaching, home blood pressure monitoring, and home-titration in controlling hypertension among low-income patients: protocol for a randomized controlled trial.' *BMC Public Health* 9: 456.
- 557 Koertke H, Zittermann A, Tenderich G, Wagner O, El-Arousy M, Krian A, Ennker J, Taborski U, Klovekorn WP, Moosdorf R, Saggau W, Koerfer R (2007). 'Low-dose oral anticoagulation in patients with mechanical heart valve prostheses: final report from the early self management anticoagulation trial II.' *Eur Heart J* 28(20): 2479-84.
- 558 Voller H, Glatz J, Taborski U, Bernardo A, Dovifat C, Heidinger K (2005). 'Self management of oral anticoagulation in nonvalvular atrial fibrillation (SMAAF study).' *Z Kardiol* 94(3): 182-6.

- 559 Christensen TD, Johnsen SP, Hjortdal VE, Hasenkam JM (2007). 'Self management of oral anticoagulant therapy: a systematic review and meta-analysis.' *Int J Cardiol* 118(1): 54-61.
- 560 Bancej CM, Campbell N, McKay DW, Nichol M, Walker RL, Kaczorowski J (2010). 'Home blood pressure monitoring among Canadian adults with hypertension: results from the 2009 Survey on Living with Chronic Diseases in Canada.' *Can J Cardiol* 26(5): e152-7.
- 561 Mailloux LU, Kapikian N, Napolitano B, et al (1996). 'Home hemodialysis: patient outcomes during a 24-year period of time from 1970 through 1993.' *Adv Ren Replace Ther* 3(2): 112-9.
- 562 Siebenhofer A, Rakovac I, Kleespies C, Piso B, Didjurgeit U (2008). 'Self management of oral anticoagulation reduces major outcomes in the elderly. A randomized controlled trial.' *Thromb Haemost* 100(6): 1089-98.
- 563 Menendez-Jandula B, Souto JC, Oliver A, Montserrat I, Quintana M, Gich I, Bonfill X, Fontcuberta J (2005). 'Comparing self management of oral anticoagulant therapy with clinic management: a randomized trial.' *Ann Intern Med* 142(1): 1-10.
- 564 Schwedes U, Siebolds M, Mertes G (2002). 'Meal-related structured self-monitoring of blood glucose: effect on diabetes control in non-insulin-treated type 2 diabetic patients.' *Diabetes Care* 25(11): 1928-32.
- 565 Guerci B, Drouin P, Grange V, et al (2003). 'Self-monitoring of blood glucose significantly improves metabolic control in patients with type 2 diabetes mellitus: the Auto-Surveillance Intervention Active (ASIA) study.' *Diabetes Metab* 29(6): 587-94.
- 566 Chase HP, Roberts MD, Wightman C, et al (2003). 'Use of the GlucoWatch biographer in children with type 1 diabetes.' *Pediatrics* 111(4): 790-4.
- 567 Sarwat S, Ilag LL, Carey MA, Shrom DS, Heine RJ (2010). 'The relationship between self-monitored blood glucose values and glycated haemoglobin in insulin-treated patients with Type 2 diabetes.' *Diabet Med* 27(5): 589-92.
- 568 Thoonen BP, Schermer TR, Van Den Boom G, et al (2003). 'Self management of asthma in general practice, asthma control and quality of life: a randomised controlled trial.' *Thorax* 58(1): 30-6.
- 569 Ignacio-Garcia JM, Gonzalez-Santos P (1995). 'Asthma self management education program by home monitoring of peak expiratory flow.' *Am J Respir Crit Care Med* 151(2 Pt 1): 353-9.
- 570 van der Meer V, van Stel HF, Bakker MJ, Roldaan AC, Assendelft WJ, Sterk PJ, Rabe KF, Sont JK (2010). 'Weekly self-monitoring and treatment adjustment benefit patients with partly controlled and uncontrolled asthma: an analysis of the SMASHING study.' *Respir Res* 11: 74.
- 571 Cappuccio FP, Kerry SM, Forbes L, Donald A (2004). 'Blood pressure control by home monitoring: meta-analysis of randomised trials.' *BMJ* 329(7458): 145.
- 572 Staessen JA, Den Hond E, Celis H, et al (2004). 'Antihypertensive treatment based on blood pressure measurement at home or in the physician's office: a randomized controlled trial.' *JAMA* 291(8): 955-64.
- 573 McCahon D, Fitzmaurice DA, Murray ET, Fuller CJ, Hobbs RF, Allan TF, Raftery JP (2003). 'SMART: self management of anticoagulation, a randomised trial.' *BMC Fam Pract* 4: 11.
- 574 Koertke H, Minami K, Boethig D, Breyman T, Seifert D, Wagner O, Atmacha N, Krian A, Ennker J, Taborski U, Klövekorn WP, Moosdorf R, Saggau W, Koerfer R (2003). 'INR self management permits lower anticoagulation levels after mechanical heart valve replacement.' *Circulation* 108(Suppl 1): I175-8.
- 575 Gardiner C, Williams K, Longair I, Mackie IJ, Machin SJ, Cohen H (2006). 'A randomised control trial of patient self management of oral anticoagulation compared with patient self-testing.' *Br J Haematol* 132(5): 598-603.
- 576 Coster S, Gulliford MC, Seed PT, et al (2000). 'Self-monitoring in type 2 diabetes mellitus: a meta-analysis.' *Diabet Med* 17(11): 755-61.
- 577 Grampian Asthma Study of Integrated Care (1994). 'Effectiveness of routine self monitoring of peak flow in patients with asthma' *BMJ* 308(6928): 564-7.
- 578 Wensley D, Silverman M (2004). 'Peak flow monitoring for guided self management in childhood asthma: a randomized controlled trial.' *Am J Respir Crit Care Med* 170(6): 606-12.
- 579 Siebenhofer A, Rakovac I, Kleespies C, Piso B, Didjurgeit U (2007). 'Self management of oral anticoagulation in the elderly: rationale, design, baselines and oral anticoagulation control after one year of follow up. A randomized controlled trial.' *Thromb Haemost* 97(3): 408-16.
- 580 Garcia-Alamino JM, Ward AM, Alonso-Coello P, Perera R, Bankhead C, Fitzmaurice D, Heneghan CJ (2010). 'Self-monitoring and self-management of oral anticoagulation.' *Cochrane Database Syst Rev* 4: CD003839.
- 581 Biermann E, Dietrich W, Standl E (2000). 'Telecare of diabetic patients with intensified insulin therapy. A randomized clinical trial.' *Stud Health Technol Inform* 77: 327-32.
- 582 Kruger DF, White K, Galpern A, et al (2003). 'Effect of modem transmission of blood glucose data on telephone consultation time, clinic work flow, and patient satisfaction for patients with gestational diabetes mellitus.' *J Am Acad Nurse Pract* 15(8): 371-5.
- 583 de Lusignan S, Wells S, Johnson P, et al (2001). 'Compliance and effectiveness of 1 year's home telemonitoring. The report of a pilot study of patients with chronic heart failure.' *Eur J Heart Fail* 3(6): 723-30.
- 584 De Clercq PA, Hasman A, Wolffenbuttel BH (2003). 'A consumer health record for supporting the patient-centered management of chronic diseases.' *Med Inform Internet Med* 28(2): 117-27.
- 585 Cho JH, Chang SA, Kwon HS, et al (2006). 'Long-term effect of the Internet-based glucose monitoring system on HbA1c reduction and glucose stability: a 30-month follow up study for diabetes management with a ubiquitous medical care system.' *Diabetes Care* 29(12): 2625-31.
- 586 Ritzema J, Troughton R, Melton I, Crozier I, Doughy R, Krum H, Walton A, Adamson P, Kar S, Shah PK, Richards M, Eigler NL, Whiting JS, Haas GJ, Heywood JT, Frampton CM, Abraham WT (2010). 'Physician-directed patient self-management of left atrial pressure in advanced chronic heart failure.' *Circulation* 121(9): 1086-95.
- 587 Balas EA, Boren SA, Griffing G (1998) . 'Computerized management of diabetes: a synthesis of controlled trials.' *Proceedings of the AMIA Annual Symposium* 295-99.
- 588 Montori VM, Helgemoe PK, Guyatt GH, et al (2004). 'Telecare for patients with type 1 diabetes and inadequate glycemic control: a randomized controlled trial and meta-analysis.' *Diabetes Care* 27(5): 1088-94.
- 589 Montani S, Bellazzi R, Quaglini S, d'Annunzio G (2001). 'Meta-analysis of the effect of the use of computer-based systems on the metabolic control of patients with diabetes mellitus.' *Diabetes Technol Ther* 3(3): 347-56.
- 590 Kwon HS, Cho JH, Kim HS, et al (2004). 'Establishment of blood glucose monitoring system using the internet.' *Diabetes Care* 27(2): 478-83.
- 591 Barlow J, Singh D, Bayer S, Curry R (2007). 'A systematic review of the benefits of home telecare for frail elderly people and those with long-term conditions.' *J Telemedicine Telecare* 13: 172-9.
- 592 McManus RJ, Mant J, Bray EP, Holder R, Jones MI, Greenfield S, Kaambwa B, Banting M, Bryan S, Little P, Williams B, Hobbs FD (2010). 'Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial.' *Lancet* 376(9736): 163-72.